

Human Slit homolog 3 protein (SLIT3) ELISA Kit

Catalog No: #EK7239



Package Size: #EK7239-1 48T #EK7239-2 96T

Orders: order@signalwayantibody.com

Support: tech@signalwayantibody.com

Description

Product Name	Human Slit homolog 3 protein (SLIT3) ELISA Kit
Brief Description	ELISA Kit
Applications	ELISA
Species Reactivity	Human (Homo sapiens)
Other Names	FLJ10764; MEGF5; SLIL2; SLIT1; Slit-3; slit2; slit homolog 3
Accession No.	O75094
Uniprot	O75094
GeneID	6586;
Storage	<p>The stability of ELISA kit is determined by the loss rate of activity. The loss rate of this kit is less than 5% within the expiration date under appropriate storage condition.</p> <p>The loss rate was determined by accelerated thermal degradation test. Keep the kit at 37C for 4 and 7 days, and compare O.D.values of the kit kept at 37C with that of at recommended temperature. (referring from China Biological Products Standard, which was calculated by the Arrhenius equation. For ELISA kit, 4 days storage at 37C can be considered as 6 months at 2 - 8C, which means 7 days at 37C equaling 12 months at 2 - 8C).</p>

Application Details

Detect Range:0.156-10 ng/mL

Sensitivity:0.071 ng/mL

Sample Type:Serum, Plasma, Other biological fluids

Sample Volume: 1-200 µL

Assay Time:1-4.5h

Detection wavelength:450 nm

Product Description

Detection Method:SandwichTest principle:This assay employs a two-site sandwich ELISA to quantitate SLIT3 in samples. An antibody specific for SLIT3 has been pre-coated onto a microplate. Standards and samples are pipetted into the wells and anySLIT3 present is bound by the immobilized antibody. After removing any unbound substances, a biotin-conjugated antibody specific for SLIT3 is added to the wells. After washing, Streptavidin conjugated Horseradish Peroxidase (HRP) is added to the wells. Following a wash to remove any unbound avidin-enzyme reagent, a substrate solution is added to the wells and color develops in proportion to the amount of SLIT3 bound in the initial step. The color development is stopped and the intensity of the color is measured.**Product Overview:**The SLIT3 cDNA encodes a 1,523-amino acid polypeptide with 41.1% similarity to the Drosophila 'slit' protein. Northern blot analysis revealed that the human SLIT3 gene was expressed as a major 5.5- and a minor 9.5-kb transcript primarily in the thyroid. SLIT1 and SLIT2 mRNAs were primarily expressed in the brain and spinal cord, respectively. In situ hybridization studies indicated that the rat Slit1 mRNA was specifically expressed in the neurons of fetal and adult forebrains. These data suggested that the SLIT genes form an evolutionarily conserved group in vertebrates and invertebrates, and that the mammalian SLIT proteins may participate in the formation and maintenance of the nervous and endocrine systems by protein-protein interactions.

Note: This product is for in vitro research use only