## Rat Heparin cofactor 2 (SERPIND1) ELISA Kit

Catalog No: #EK7371

SAB Signalway Antibody

Orders: order@signalwayantibody.com Support: tech@signalwayantibody.com

Package Size: #EK7371-1 48T #EK7371-2 96T

Description	
Product Name	Rat Heparin cofactor 2 (SERPIND1) ELISA Kit
Brief Description	ELISA Kit
Applications	ELISA
Species Reactivity	Rat (Rattus norvegicus)
Other Names	D22S673; HC2; HCF2; HCII; HLS2; LS2; OTTHUMP00000198451 heparin cofactor II leuserpin 2 serine (or
	cysteine) proteinase inhibitor; clade D (heparin cofactor); member 1
Accession No.	P47776
Uniprot	P47776
GeneID	100009170;
Storage	The stability of ELISA kit is determined by the loss rate of activity. The loss rate of this kit is less than 5%
	within the expiration date under appropriate storage condition.
	The loss rate was determined by accelerated thermal degradation test. Keep the kit at 37C for 4 and 7 days,
	and compare O.D.values of the kit kept at 37C with that of at recommended temperature. (referring from China
	Biological Products Standard, which was calculated by the Arrhenius equation. For ELISA kit, 4 days storage
	at 37C can be considered as 6 months at 2 - 8C, which means 7 days at 37C equaling 12 months at 2 - 8C).
	The loss rate was determined by accelerated thermal degradation test. Keep the kit at 37C for 4 and 7 days, and compare O.D.values of the kit kept at 37C with that of at recommended temperature. (referring from China Biological Products Standard, which was calculated by the Arrhenius equation. For ELISA kit, 4 days storage

## **Application Details**

Detect Range:7.81-500 ng/mL	
Sensitivity:3.3 ng/mL	
Sample Type:Serum, Plasma, Other biological fluids	
Sample Volume: 1-200 μL	
Assay Time:1-4.5h	
Detection wavelength:450 nm	

## **Product Description**

Detection Method:SandwichTest principle:This assay employs a two-site sandwich ELISA to quantitate SERPIND1 in samples. An antibody specific for SERPIND1 has been pre-coated onto a microplate. Standards and samples are pipetted into the wells and anySERPIND1 present is bound by the immobilized antibody. After removing any unbound substances, a biotin-conjugated antibody specific for SERPIND1 is added to the wells. After washing, Streptavidin conjugated Horseradish Peroxidase (HRP) is added to the wells. Following a wash to remove any unbound avidin-enzyme reagent, a substrate solution is added to the wells and color develops in proportion to the amount of SERPIND1 bound in the initial step. The color development is stopped and the intensity of the color is measured. Product Overview: Heparin cofactor II, a protein encoded by the SERPIND1 gene, is a coagulation factor that inhibits IIa, and is a cofactor for heparin and dermatan sulfate ("minor antithrombin").

The product encoded by this gene is a serine proteinase inhibitor which rapidly inhibits thrombin in the presence of dermatan sulfate or heparin. The gene contains five exons and four introns.

This protein shares homology with antithrombin III and other members of the alpha 1-antitrypsin superfamily. Mutations in this gene are associated with heparin cofactor II deficiency. Heparin Cofactor II deficiency can lead to increased thrombin generation and a hypercoagulable state.

Note: This product is for in vitro research use only