

Human Secretogranin-2 (SCG2) ELISA Kit

Catalog No: #EK7441



Package Size: #EK7441-1 48T #EK7441-2 96T

Orders: order@signalwayantibody.com

Support: tech@signalwayantibody.com

Description

Product Name	Human Secretogranin-2 (SCG2) ELISA Kit
Brief Description	ELISA Kit
Applications	ELISA
Species Reactivity	Human (Homo sapiens)
Other Names	CHGC; SN; SgII; Chromogranin C (secretogranin II) EM66 secretogranin II secretoneurin
Accession No.	P13521
Uniprot	P13521
GeneID	7857;
Storage	<p>The stability of ELISA kit is determined by the loss rate of activity. The loss rate of this kit is less than 5% within the expiration date under appropriate storage condition.</p> <p>The loss rate was determined by accelerated thermal degradation test. Keep the kit at 37C for 4 and 7 days, and compare O.D.values of the kit kept at 37C with that of at recommended temperature. (referring from China Biological Products Standard, which was calculated by the Arrhenius equation. For ELISA kit, 4 days storage at 37C can be considered as 6 months at 2 - 8C, which means 7 days at 37C equaling 12 months at 2 - 8C).</p>

Application Details

Detect Range:1.25-80 ng/mL

Sensitivity:0.49 ng/mL

Sample Type:Serum, Plasma, Other biological fluids

Sample Volume: 1-200 µL

Assay Time:1-4.5h

Detection wavelength:450 nm

Product Description

Detection Method:SandwichTest principle:This assay employs a two-site sandwich ELISA to quantitate SCG2 in samples. An antibody specific for SCG2 has been pre-coated onto a microplate. Standards and samples are pipetted into the wells and anySCG2 present is bound by the immobilized antibody. After removing any unbound substances, a biotin-conjugated antibody specific for SCG2 is added to the wells. After washing, Streptavidin conjugated Horseradish Peroxidase (HRP) is added to the wells. Following a wash to remove any unbound avidin-enzyme reagent, a substrate solution is added to the wells and color develops in proportion to the amount of SCG2 bound in the initial step. The color development is stopped and the intensity of the color is measured.**Product Overview:**SCG2 is a member of the chromogranin/secretogranin family of neuroendocrine secretory proteins. Studies in rodents suggest that the full-length protein, secretogranin II, is involved in the packaging or sorting of peptide hormones and neuropeptides into secretory vesicles. The full-length protein is cleaved to produce the active peptide secretoneurin, which exerts chemotaxic effects on specific cell types, and EM66, whose function is unknown.SC2 also has 9 potential dibasic cleavage sites, an N-glycosylation site not found in the bovine protein, and a putative sulfation site (tyr121). SC2 was predicted to assume a secondary structure of alternating helix and turn structures over 45% and 40% of its length, respectively. The authors suggested that this structure may coordinate calcium.

Note: This product is for in vitro research use only