

Rat ATP-binding cassette sub-family G member 2 (ABCG2) ELISA Kit



Catalog No: #EK7629

Orders: order@signalwayantibody.com

Package Size: #EK7629-1 48T #EK7629-2 96T

Support: tech@signalwayantibody.com

Description

Product Name	Rat ATP-binding cassette sub-family G member 2 (ABCG2) ELISA Kit
Brief Description	ELISA Kit
Applications	ELISA
Species Reactivity	Rat (<i>Rattus norvegicus</i>)
Other Names	ABC15; ABCP; BCRP; BCRP1; BMDP; CD338; CDw338; EST157481; MGC102821; MRX; MXR; MXR1; ABC transporter ATP-binding cassette transporter G2 ATP-binding cassette; sub-family G; member 2 breast cancer re
Accession No.	Q80W57
Uniprot	Q80W57
GeneID	312382;
Storage	The stability of ELISA kit is determined by the loss rate of activity. The loss rate of this kit is less than 5% within the expiration date under appropriate storage condition. The loss rate was determined by accelerated thermal degradation test. Keep the kit at 37C for 4 and 7 days, and compare O.D.values of the kit kept at 37C with that of at recommended temperature. (referring from China Biological Products Standard, which was calculated by the Arrhenius equation. For ELISA kit, 4 days storage at 37C can be considered as 6 months at 2 - 8C, which means 7 days at 37C equaling 12 months at 2 - 8C).

Application Details

Detect Range:0.312-20 ng/mL

Sensitivity:0.128 ng/mL

Sample Type:Serum, Plasma, Other biological fluids

Sample Volume: 1-200 µL

Assay Time:1-4.5h

Detection wavelength:450 nm

Product Description

Detection Method:SandwichTest principle:This assay employs a two-site sandwich ELISA to quantitate ABCG2 in samples. An antibody specific for ABCG2 has been pre-coated onto a microplate. Standards and samples are pipetted into the wells and anyABCG2 present is bound by the immobilized antibody. After removing any unbound substances, a biotin-conjugated antibody specific for ABCG2 is added to the wells. After washing, Streptavidin conjugated Horseradish Peroxidase (HRP) is added to the wells. Following a wash to remove any unbound avidin-enzyme reagent, a substrate solution is added to the wells and color develops in proportion to the amount of ABCG2 bound in the initial step. The color development is stopped and the intensity of the color is measured.**Product Overview:**ATP-binding cassette, sub-family G (WHITE), member 2 is included in the superfamily of ATP-binding cassette (ABC) transporters. ABC proteins transport various molecules across extra- and intra-cellular membranes. ABC genes are divided into seven distinct subfamilies (ABC1, MDR/TAP, MRP, ALD, OABP, GCN20, White). This protein is a member of the White subfamily. Alternatively referred to as a breast cancer resistance protein, this protein functions as a xenobiotic transporter which may play a major role in multi-drug resistance. It likely serves as a cellular defense mechanism in response to mitoxantrone and anthracycline exposure. Significant

expression of this protein has been observed in the placenta, which may suggest a potential role for this molecule in placenta tissue.

Note: This product is for in vitro research use only