Pig Angiotensin Converting Enzyme (ACE) ELISA Kit

Catalog No: #EK7644



Package Size: #EK7644-1 48T #EK7644-2 96T

Orders: order@signalwayantibody.com Support: tech@signalwayantibody.com

Description	
Product Name	Pig Angiotensin Converting Enzyme (ACE) ELISA Kit
Brief Description	ELISA Kit
Applications	ELISA
Species Reactivity	Pig (Sus scrofa; Porcine)
Other Names	ACE1; CD143; DCP; DCP1; MGC26566; MVCD3; CD143 antigen angiotensin I converting enzyme
	1 angiotensin converting enzyme; somatic isoform carboxycathepsin dipeptidyl carboxypeptidase 1 kininase
	II pep
Accession No.	P12821
Uniprot	P12821
GeneID	1636;
Storage	The stability of ELISA kit is determined by the loss rate of activity. The loss rate of this kit is less than 5%
	within the expiration date under appropriate storage condition.
	The loss rate was determined by accelerated thermal degradation test. Keep the kit at 37C for 4 and 7 days,
	and compare O.D.values of the kit kept at 37C with that of at recommended temperature. (referring from China
	Biological Products Standard, which was calculated by the Arrhenius equation. For ELISA kit, 4 days storage
	at 37C can be considered as 6 months at 2 - 8C, which means 7 days at 37C equaling 12 months at 2 - 8C).

Application Details

Detect Range:31.25-2000 pg/mL
Sensitivity:11.2 pg/mL
Sample Type:Serum, Plasma, Other biological fluids
Sample Volume: 1-200 μL
Assay Time:1-4.5h
Detection wavelength:450 nm

Product Description

Detection Method:SandwichTest principle:This assay employs a two-site sandwich ELISA to quantitate ACE in samples. An antibody specific for ACE has been pre-coated onto a microplate. Standards and samples are pipetted into the wells and anyACE present is bound by the immobilized antibody. After removing any unbound substances, a biotin-conjugated antibody specific for ACE is added to the wells. After washing, Streptavidin conjugated Horseradish Peroxidase (HRP) is added to the wells. Following a wash to remove any unbound avidin-enzyme reagent, a substrate solution is added to the wells and color develops in proportion to the amount of ACE bound in the initial step. The color development is stopped and the intensity of the color is measured. Product Overview: Angiotensin I-converting enzyme (ACE, EC 3.4.15.1), an exopeptidase, is a circulating enzyme that participates in the body's renin-angiotensin system (RAS), which mediates extracellular volume (i.e. that of the blood plasma, lymph and interstitial fluid), and arterial vasoconstriction. It is secreted by pulmonary and renal endothelial cells and catalyzes the conversion of decapeptide angiotensin I to octapeptide angiotensin II. ACE Ubiquitously expressed, with highest levels in lung, kidney, heart, gastrointestinal system and prostate. The testis-specific isoforn is expressed in spermatocytes, adult testis. ACE can Convert angiotensin I to angiotensin II by release of the terminal His-Leu, this results in an increase of the vasoconstrictor activity of angiotensin.

Note: This product is for in vitro research use only