## **Product Datasheet**

## Human Anterior gradient protein 2 homolog (AGR2) ELISA Kit

Catalog No: #EK7787

Package Size: #EK7787-1 48T #EK7787-2 96T



Orders: order@signalwayantibody.com Support: tech@signalwayantibody.com

Description		
Product Name	Human Anterior gradient protein 2 homolog (AGR2) ELISA Kit	
Brief Description	ELISA Kit	
Applications	ELISA	
Species Reactivity	Human (Homo sapiens)	
Other Names	AG2; GOB-4; HAG-2; PDIA17; XAG-2; anterior gradient 2 homolog protein disulfide isomerase family A;	
	member 17 secreted cement gland homolog	
Accession No.	O95994	
Uniprot	O95994	
GeneID	10551;	
Storage	The stability of ELISA kit is determined by the loss rate of activity. The loss rate of this kit is less than 5%	
	within the expiration date under appropriate storage condition.	
	The loss rate was determined by accelerated thermal degradation test. Keep the kit at 37C for 4 and 7 days,	
	and compare O.D.values of the kit kept at 37C with that of at recommended temperature. (referring from China	
	Biological Products Standard, which was calculated by the Arrhenius equation. For ELISA kit, 4 days storage	
	at 37C can be considered as 6 months at 2 - 8C, which means 7 days at 37C equaling 12 months at 2 - 8C).	

## **Application Details**

Detect Range:1.56-100 ng/mL	
Sensitivity:0.62 ng/mL	
Sample Type:Serum, Plasma, Other biological fluids	
Sample Volume: 1-200 µL	
Assay Time:1-4.5h	
Detection wavelength:450 nm	

## **Product Description**

Detection Method:SandwichTest principle:This assay employs a two-site sandwich ELISA to quantitate AGR2 in samples. An antibody specific for AGR2 has been pre-coated onto a microplate. Standards and samples are pipetted into the wells and anyAGR2 present is bound by the immobilized antibody. After removing any unbound substances, a biotin-conjugated antibody specific for AGR2 is added to the wells. After washing, Streptavidin conjugated Horseradish Peroxidase (HRP) is added to the wells. Following a wash to remove any unbound avidin-enzyme reagent, a substrate solution is added to the wells and color develops in proportion to the amount of AGR2 bound in the initial step. The color development is stopped and the intensity of the color is measured.Product Overview:Anterior gradient homolog 2 (AGR2) is a human protein originally discovered in Xenopus laevis, encoded for by the gene of the same name. In Xenopus AGR2 plays a role in cement gland differentiation, however in human cancer cell lines high levels of AGR2 correlate with downregulation of the p53 response and AGR2 levels are elevated in the preneoplastic tissue Barrett's oesophagus. The deduced 175-amino acid soluble AGR2 protein, which is 91% identical to the mouse protein and 47% identical to the frog protein, contains a signal peptide. Northern blot analysis revealed strongest expression of 0.9- and 1.6-kb AGR2 transcripts in lung and in all ER-positive breast carcinoma lines tested; weaker expression was detected in pancreas.

Note: This product is for in vitro research use only