

Bovine Pyrroline-5-carboxylate reductase 3 (PYCRL) ELISA Kit



Catalog No: #EK7902

Orders: order@signalwayantibody.com

Package Size: #EK7902-1 48T #EK7902-2 96T

Support: tech@signalwayantibody.com

Description

Product Name	Bovine Pyrroline-5-carboxylate reductase 3 (PYCRL) ELISA Kit
Brief Description	ELISA Kit
Applications	ELISA
Species Reactivity	Bovine (Bos taurus; Cattle)
Other Names	FLJ13852;
Accession No.	Q58D08
Uniprot	Q58D08
GeneID	512526;
Storage	<p>The stability of ELISA kit is determined by the loss rate of activity. The loss rate of this kit is less than 5% within the expiration date under appropriate storage condition.</p> <p>The loss rate was determined by accelerated thermal degradation test. Keep the kit at 37C for 4 and 7 days, and compare O.D.values of the kit kept at 37C with that of at recommended temperature. (referring from China Biological Products Standard, which was calculated by the Arrhenius equation. For ELISA kit, 4 days storage at 37C can be considered as 6 months at 2 - 8C, which means 7 days at 37C equaling 12 months at 2 - 8C).</p>

Application Details

Detect Range:Request Information

Sensitivity:Request Information

Sample Type:Serum, Plasma, Other biological fluids

Sample Volume: 1-200 µL

Assay Time:1-4.5h

Detection wavelength:450 nm

Product Description

Detection Method:SandwichTest principle:This assay employs a two-site sandwich ELISA to quantitate PYCRL in samples. An antibody specific for PYCRL has been pre-coated onto a microplate. Standards and samples are pipetted into the wells and anyPYCRL present is bound by the immobilized antibody. After removing any unbound substances, a biotin-conjugated antibody specific for PYCRL is added to the wells. After washing, Streptavidin conjugated Horseradish Peroxidase (HRP) is added to the wells. Following a wash to remove any unbound avidin-enzyme reagent, a substrate solution is added to the wells and color develops in proportion to the amount of PYCRL bound in the initial step. The color development is stopped and the intensity of the color is measured.Product Overview:PYCRL belongs to the pyrroline-5-carboxylate reductase family.

Pyrroline-5-carboxylate reductase catalyzes the the NAD(P)H-dependent conversion of pyrroline-5-carboxylate to proline. Pyrroline-5-carboxylate reductase(P5CR) is the enzyme that catalyzes the terminal step in the biosynthesis of proline from glutamate, the NAD(P) dependent oxidation of 1-pyrroline-5-carboxylate into proline.

Pyrroline-5-carboxylate reductase, which converts pyrroline-5-carboxylate to proline, has been identified in human erythrocytes. The level of pyrroline-5-carboxylate reductase activity in these cells is comparable to the activity levels of major erythrocyte enzymes. The physiologic function of the enzyme in erythrocytes cannot be related to its function in other tissues.

Note: This product is for in vitro research use only