Mouse Polyserase-2 (PRSS36) ELISA Kit

Catalog No: #EK8083

Package Size: #EK8083-1 48T #EK8083-2 96T



Orders: order@signalwayantibody.com Support: tech@signalwayantibody.com

| Description | |
|--------------------|--|
| Product Name | Mouse Polyserase-2 (PRSS36) ELISA Kit |
| Brief Description | ELISA Kit |
| Applications | ELISA |
| Species Reactivity | Mouse (Mus musculus) |
| Other Names | FLJ90661; polyserase-2 polyserine protease-2 |
| Accession No. | Q5K2P8 |
| Uniprot | Q5K2P8 |
| Storage | The stability of ELISA kit is determined by the loss rate of activity. The loss rate of this kit is less than 5% |

within the expiration date under appropriate storage condition.

The loss rate was determined by accelerated thermal degradation test. Keep the kit at 37C for 4 and 7 days, and compare O.D.values of the kit kept at 37C with that of at recommended temperature. (referring from China Biological Products Standard, which was calculated by the Arrhenius equation. For ELISA kit, 4 days storage

at 37C can be considered as 6 months at 2 - 8C, which means 7 days at 37C equaling 12 months at 2 - 8C).

Application Details

| Detect Range:Request Information |
|--|
| Sensitivity:Request Information |
| Sample Type:Serum, Plasma, Other biological fluids |
| Sample Volume: 1-200 μL |
| Assay Time:1-4.5h |
| Detection wavelength:450 nm |

Product Description

Detection Method:SandwichTest principle:This assay employs a two-site sandwich ELISA to quantitate PRSS36 in samples. An antibody specific for PRSS36 has been pre-coated onto a microplate. Standards and samples are pipetted into the wells and anyPRSS36 present is bound by the immobilized antibody. After removing any unbound substances, a biotin-conjugated antibody specific for PRSS36 is added to the wells. After washing, Streptavidin conjugated Horseradish Peroxidase (HRP) is added to the wells. Following a wash to remove any unbound avidin-enzyme reagent, a substrate solution is added to the wells and color develops in proportion to the amount of PRSS36 bound in the initial step. The color development is stopped and the intensity of the color is measured. Product Overview: The deduced 855-amino acid protein has a predicted molecular mass of 93 kD and contains 9 potential N-glycosylation sites. PRSS36 contains a signal peptide, followed by a propeptide domain, an active serine protease domain with the catalytic his, asp, and ser residues, and 2 serine protease domains predicted to be catalytically inactive. The first serine protease domain shares amino acid identity of 40 to 42% with gamma-1 tryptase (TPSG1), pancreasin, matriptase (ST14;), matriptase-2 (TMPRSS6), and polyserase-1 (TMPRSS9). Northern blot analysis detected a 5-kb transcript in fetal kidney and adult skeletal muscle, liver, placenta, and heart. A 2.2-kb transcript was detected in placenta, prostate, and colon and lung adenocarcinoma cell lines, and may be produced by alternative splicing.

Note: This product is for in vitro research use only