

Mouse Protein kinase C epsilon type (PRKCE) ELISA Kit



Catalog No: #EK8242

Orders: order@signalwayantibody.com

Package Size: #EK8242-1 48T #EK8242-2 96T

Support: tech@signalwayantibody.com

Description

Product Name	Mouse Protein kinase C epsilon type (PRKCE) ELISA Kit
Brief Description	ELISA Kit
Applications	ELISA
Species Reactivity	Mouse (Mus musculus)
Other Names	MGC125656; MGC125657; PKCE; nPKC-epsilon;
Accession No.	P16054
Uniprot	P16054
GeneID	18754;
Storage	<p>The stability of ELISA kit is determined by the loss rate of activity. The loss rate of this kit is less than 5% within the expiration date under appropriate storage condition.</p> <p>The loss rate was determined by accelerated thermal degradation test. Keep the kit at 37C for 4 and 7 days, and compare O.D.values of the kit kept at 37C with that of at recommended temperature. (referring from China Biological Products Standard, which was calculated by the Arrhenius equation. For ELISA kit, 4 days storage at 37C can be considered as 6 months at 2 - 8C, which means 7 days at 37C equaling 12 months at 2 - 8C).</p>

Application Details

Detect Range:1.56-100 ng/mL

Sensitivity:0.61 ng/mL

Sample Type:Serum, Plasma, Other biological fluids

Sample Volume: 1-200 µL

Assay Time:1-4.5h

Detection wavelength:450 nm

Product Description

Detection Method:SandwichTest principle:This assay employs a two-site sandwich ELISA to quantitate PRKCE in samples. An antibody specific for PRKCE has been pre-coated onto a microplate. Standards and samples are pipetted into the wells and anyPRKCE present is bound by the immobilized antibody. After removing any unbound substances, a biotin-conjugated antibody specific for PRKCE is added to the wells. After washing, Streptavidin conjugated Horseradish Peroxidase (HRP) is added to the wells. Following a wash to remove any unbound avidin-enzyme reagent, a substrate solution is added to the wells and color develops in proportion to the amount of PRKCE bound in the initial step. The color development is stopped and the intensity of the color is measured.**Product Overview:**PKC family members also serve as major receptors for phorbol esters, a class of tumor promoters. Each member of the PKC family has a specific expression profile and is believed to play a distinct role in cells. The protein encoded by this gene is one of the PKC family members.

Protein kinase C epsilon type kinase has been shown to be involved in many different cellular functions, such as neuron channel activation, apoptosis, cardioprotection from ischemia, heat shock response, as well as insulin exocytosis. Knockout studies in mice suggest that this kinase is important for lipopolysaccharide (LPS)-mediated signaling in activated macrophages and may also play a role in controlling anxiety-like behavior.

Note: This product is for in vitro research use only