

Human Putative phospholipase B-like 2 (PLBD2) ELISA Kit

Catalog No: #EK8450

Orders: order@signalwayantibody.com

Package Size: #EK8450-1 48T #EK8450-2 96T

Support: tech@signalwayantibody.com

Description

Product Name	Human Putative phospholipase B-like 2 (PLBD2) ELISA Kit
Brief Description	ELISA Kit
Applications	ELISA
Species Reactivity	Human (Homo sapiens)
Other Names	P76; LAMA-like protein 2 PLB homolog 2 lamina ancestor homolog 2 mannose-6-phosphate protein associated protein p76 phospholipase B-like 2 32 kDa form phospholipase B-like 2 45 kDa form
Accession No.	Q2KIY5
Uniprot	Q2KIY5
GeneID	514347;
Storage	The stability of ELISA kit is determined by the loss rate of activity. The loss rate of this kit is less than 5% within the expiration date under appropriate storage condition. The loss rate was determined by accelerated thermal degradation test. Keep the kit at 37C for 4 and 7 days, and compare O.D.values of the kit kept at 37C with that of at recommended temperature. (referring from China Biological Products Standard, which was calculated by the Arrhenius equation. For ELISA kit, 4 days storage at 37C can be considered as 6 months at 2 - 8C, which means 7 days at 37C equaling 12 months at 2 - 8C).

Application Details

Detect Range:78.1-5000 pg/mL

Sensitivity:35 pg/mL

Sample Type:Serum, Plasma, Other biological fluids

Sample Volume: 1-200 µL

Assay Time:1-4.5h

Detection wavelength:450 nm

Product Description

Detection Method:Sandwich Test principle:This assay employs a two-site sandwich ELISA to quantitate PLBD2 in samples. An antibody specific for PLBD2 has been pre-coated onto a microplate. Standards and samples are pipetted into the wells and any PLBD2 present is bound by the immobilized antibody. After removing any unbound substances, a biotin-conjugated antibody specific for PLBD2 is added to the wells. After washing, Streptavidin conjugated Horseradish Peroxidase (HRP) is added to the wells. Following a wash to remove any unbound avidin-enzyme reagent, a substrate solution is added to the wells and color develops in proportion to the amount of PLBD2 bound in the initial step. The color development is stopped and the intensity of the color is measured.

Product Overview:Most soluble lysosomal proteins carry Man6P (mannose 6-phosphate), a specific carbohydrate marker that enables their binding to cellular MPRs (Man6P receptors) and their subsequent targeting towards the lysosome.

This characteristic was exploited to identify novel soluble lysosomal proteins by proteomic analysis of Man6P proteins purified from a human cell line. The presence of Man6P sugars was confirmed by an MPR overlay experiment, which showed the direct and Man6P-dependent interaction between p76 and the MPR. The presence of six N-glycosylation sites was validated by progressive peptide-N-glycosidase F deglycosylation. Experiments using N- and C-termini directed anti-p76 antibodies provided insights into p76 maturation.

Note: This product is for in vitro research use only