

Rat Polymeric immunoglobulin receptor (PIGR) ELISA Kit



Catalog No: #EK8514

Orders: order@signalwayantibody.com

Package Size: #EK8514-1 48T #EK8514-2 96T

Support: tech@signalwayantibody.com

Description

Product Name	Rat Polymeric immunoglobulin receptor (PIGR) ELISA Kit
Brief Description	ELISA Kit
Applications	ELISA
Species Reactivity	Rat (<i>Rattus norvegicus</i>)
Other Names	FLJ22667; MGC125361; MGC125362; hepatocellular carcinoma associated protein TB6
Accession No.	P01832
Uniprot	P01832
GeneID	100328593;
Storage	<p>The stability of ELISA kit is determined by the loss rate of activity. The loss rate of this kit is less than 5% within the expiration date under appropriate storage condition.</p> <p>The loss rate was determined by accelerated thermal degradation test. Keep the kit at 37C for 4 and 7 days, and compare O.D.values of the kit kept at 37C with that of at recommended temperature. (referring from China Biological Products Standard, which was calculated by the Arrhenius equation. For ELISA kit, 4 days storage at 37C can be considered as 6 months at 2 - 8C, which means 7 days at 37C equaling 12 months at 2 - 8C).</p>

Application Details

Detect Range:31.25-2000 ng/mL

Sensitivity:7.8 ng/mL

Sample Type:Serum, Plasma, Other biological fluids

Sample Volume: 1-200 μ L

Assay Time:1-4.5h

Detection wavelength:450 nm

Product Description

Detection Method:SandwichTest principle:This assay employs a two-site sandwich ELISA to quantitate PIGR in samples. An antibody specific for PIGR has been pre-coated onto a microplate. Standards and samples are pipetted into the wells and anyPIGR present is bound by the immobilized antibody. After removing any unbound substances, a biotin-conjugated antibody specific for PIGR is added to the wells. After washing, Streptavidin conjugated Horseradish Peroxidase (HRP) is added to the wells. Following a wash to remove any unbound avidin-enzyme reagent, a substrate solution is added to the wells and color develops in proportion to the amount of PIGR bound in the initial step. The color development is stopped and the intensity of the color is measured.**Product Overview:**Polymeric immunoglobulin receptor is a Fc receptor which facilitates the secretion of IgA and IgM. The poly-Ig receptor is expressed on several glandular epithelia including those of liver and breast. It mediates transcellular transport of polymeric immunoglobulin molecules.

It is a member of the immunoglobulin superfamily (Hood et al., 1985). The receptor has 5 units with homology to the variable (V) units of immunoglobulins and a transmembrane region, which also has some homology to certain immunoglobulin variable regions.

Note: This product is for in vitro research use only