## Human 11-Deoxycortisol ELISA Kit

Catalog No: #EK8730

Package Size: #EK8730-1 48T #EK8730-2 96T



Orders: order@signalwayantibody.com Support: tech@signalwayantibody.com

## Description

| Product Name       | Human 11-Deoxycortisol ELISA Kit   |  |  |
|--------------------|--|--|--|
| Brief Description  | ELISA Kit  |  |  |
| Applications       | ELISA  |  |  |
| Species Reactivity | Human (Homo sapiens)   |  |  |
| Storage            | The stability of ELISA kit is determined by the loss rate of activity. The loss rate of this kit is less than 5% |  |  |
|                    | within the expiration date under appropriate storage condition.  |  |  |
|                    | The loss rate was determined by accelerated thermal degradation test. Keep the kit at 37C for 4 and 7 days,      |  |  |
|                    | and compare O.D.values of the kit kept at 37C with that of at recommended temperature. (referring from China     |  |  |
|                    | Biological Products Standard, which was calculated by the Arrhenius equation. For ELISA kit, 4 days storage      |  |  |
|                    | at 37C can be considered as 6 months at 2 - 8C, which means 7 days at 37C equaling 12 months at 2 - 8C).         |  |  |
|                    |  |  |  |

| Application Details          |                        |  |
|------------------------------|------------------------|--|
| Detect Range:0.31-20 pmol/mL |                        |  |
| Sensitivity:0.1 pmol/mL      |                        |  |
| Sample Type:Serum, Plasma, C | ther biological fluids |  |
| Sample Volume: 1-200 µL      |                        |  |
| Assay Time:1-4.5h            |                        |  |
| Detection wavelength:450 nm  |                        |  |

## **Product Description**

Detection Method:SandwichTest principle:This assay employs a two-site sandwich ELISA to quantitate 11-Deoxycortisol in samples. An antibody specific for 11-Deoxycortisol has been pre-coated onto a microplate. Standards and samples are pipetted into the wells and any11-Deoxycortisol present is bound by the immobilized antibody. After removing any unbound substances, a biotin-conjugated antibody specific for 11-Deoxycortisol is added to the wells. After washing, Streptavidin conjugated Horseradish Peroxidase (HRP) is added to the wells. Following a wash to remove any unbound avidin-enzyme reagent, a substrate solution is added to the wells and color develops in proportion to the amount of 11-Deoxycortisol bound in the initial step. The color development is stopped and the intensity of the color is measured.

Note: This product is for in vitro research use only