## Deer Urateoxidase ELISA Kit

Catalog No: #EK8735

Package Size: #EK8735-1 48T #EK8735-2 96T



Orders: order@signalwayantibody.com Support: tech@signalwayantibody.com

D	esc	cri	pt	io	n

Product Name	Deer Urateoxidase ELISA Kit
Brief Description	ELISA Kit
Applications	ELISA
Species Reactivity	Deer
Storage	The stability of ELISA kit is determined by the loss rate of activity. The loss rate of this kit is less than 5%
	within the expiration date under appropriate storage condition.
	The loss rate was determined by accelerated thermal degradation test. Keep the kit at 37C for 4 and 7 days,
	and compare O.D.values of the kit kept at 37C with that of at recommended temperature. (referring from China
	Biological Products Standard, which was calculated by the Arrhenius equation. For ELISA kit, 4 days storage
	at 37C can be considered as 6 months at 2 - 8C, which means 7 days at 37C equaling 12 months at 2 - 8C).

## **Application Details**

Detect Range:31.25-2000 pg/mL
Sensitivity:15 pg/mL
Sample Type:Serum, Plasma, Other biological fluids
Sample Volume: 1-200 μL
Assay Time:1-4.5h
Detection wavelength:450 nm

## **Product Description**

Detection Method:SandwichTest principle:This assay employs a two-site sandwich ELISA to quantitate Urateoxidase in samples. An antibody specific for Urateoxidase has been pre-coated onto a microplate. Standards and samples are pipetted into the wells and anyUrateoxidase present is bound by the immobilized antibody. After removing any unbound substances, a biotin-conjugated antibody specific for Urateoxidase is added to the wells. After washing, Streptavidin conjugated Horseradish Peroxidase (HRP) is added to the wells. Following a wash to remove any unbound avidin-enzyme reagent, a substrate solution is added to the wells and color develops in proportion to the amount of Urateoxidase bound in the initial step. The color development is stopped and the intensity of the color is measured.

Note: This product is for in vitro research use only