Human Nanos homolog 2 (NANOS2) ELISA Kit

Catalog No: #EK8859

Description

SAB Signalway Antibody

Package Size: #EK8859-1 48T #EK8859-2 96T

Orders: order@signalwayantibody.com Support: tech@signalwayantibody.com

Brief Description ELISA Kit Applications ELISA Species Reactivity Human (Homo sapiens) Other Names NOS2; nanos homolog 2 Accession No. P60321 Uniprot P60321 GeneID 339345;	Bootinplion	
Applications ELISA Species Reactivity Human (Homo sapiens) Other Names NOS2; nanos homolog 2 Accession No. P60321 Uniprot P60321 GeneID 339345;	Product Name	Human Nanos homolog 2 (NANOS2) ELISA Kit
Species Reactivity Human (Homo sapiens) Other Names NOS2; nanos homolog 2 Accession No. P60321 Uniprot P60321 GeneID 339345;	Brief Description	ELISA Kit
Other Names NOS2; nanos homolog 2 Accession No. P60321 Uniprot P60321 GeneID 339345;	Applications	ELISA
Accession No. P60321 Uniprot P60321 GeneID 339345;	Species Reactivity	Human (Homo sapiens)
Uniprot P60321 GeneID 339345;	Other Names	NOS2; nanos homolog 2
GeneID 339345;	Accession No.	P60321
	Uniprot	P60321
Storage The stability of ELISA kit is determined by the loss rate of activity. The loss rate of this kit is less than 5%	GeneID	339345;
	Storage	The stability of ELISA kit is determined by the loss rate of activity. The loss rate of this kit is less than 5%

within the expiration date under appropriate storage condition.

The loss rate was determined by accelerated thermal degradation test. Keep the kit at 37C for 4 and 7 days, and compare O.D.values of the kit kept at 37C with that of at recommended temperature. (referring from China Biological Products Standard, which was calculated by the Arrhenius equation. For ELISA kit, 4 days storage at 37C can be considered as 6 months at 2 - 8C, which means 7 days at 37C equaling 12 months at 2 - 8C).

Application Details

Detect Range:Request Information		
Sensitivity:Request Information		
Sample Type:Serum, Plasma, Other biological fluids		
Sample Volume: 1-200 μL		
Assay Time:1-4.5h		
Detection wavelength:450 nm		

Product Description

Detection Method:SandwichTest principle:This assay employs a two-site sandwich ELISA to quantitate NANOS2 in samples. An antibody specific for NANOS2 has been pre-coated onto a microplate. Standards and samples are pipetted into the wells and anyNANOS2 present is bound by the immobilized antibody. After removing any unbound substances, a biotin-conjugated antibody specific for NANOS2 is added to the wells. After washing, Streptavidin conjugated Horseradish Peroxidase (HRP) is added to the wells. Following a wash to remove any unbound avidin-enzyme reagent, a substrate solution is added to the wells and color develops in proportion to the amount of NANOS2 bound in the initial step. The color development is stopped and the intensity of the color is measured.

Note: This product is for in vitro research use only