

## Rat Myocilin (MYOC) ELISA Kit

Catalog No: #EK8950



Package Size: #EK8950-1 48T #EK8950-2 96T

Orders: [order@signalwayantibody.com](mailto:order@signalwayantibody.com)Support: [tech@signalwayantibody.com](mailto:tech@signalwayantibody.com)

## Description

Product Name	Rat Myocilin (MYOC) ELISA Kit
Brief Description	ELISA Kit
Applications	ELISA
Species Reactivity	Rat ( <i>Rattus norvegicus</i> )
Other Names	GLC1A; GPOA; JOAG; JOAG1; TIGR; myocilin; mutated trabecular meshwork-induced glucocorticoid response protein myocilin
Accession No.	Q866N2
Uniprot	Q866N2
GeneID	100008897;
Storage	<p>The stability of ELISA kit is determined by the loss rate of activity. The loss rate of this kit is less than 5% within the expiration date under appropriate storage condition.</p> <p>The loss rate was determined by accelerated thermal degradation test. Keep the kit at 37C for 4 and 7 days, and compare O.D.values of the kit kept at 37C with that of at recommended temperature. (referring from China Biological Products Standard, which was calculated by the Arrhenius equation. For ELISA kit, 4 days storage at 37C can be considered as 6 months at 2 - 8C, which means 7 days at 37C equaling 12 months at 2 - 8C).</p>

## Application Details

Detect Range:Request Information

Sensitivity:Request Information

Sample Type:Serum, Plasma, Other biological fluids

Sample Volume: 1-200 µL

Assay Time:1-4.5h

Detection wavelength:450 nm

## Product Description

Detection Method:SandwichTest principle:This assay employs a two-site sandwich ELISA to quantitate MYOC in samples. An antibody specific for MYOC has been pre-coated onto a microplate. Standards and samples are pipetted into the wells and anyMYOC present is bound by the immobilized antibody. After removing any unbound substances, a biotin-conjugated antibody specific for MYOC is added to the wells. After washing, Streptavidin conjugated Horseradish Peroxidase (HRP) is added to the wells. Following a wash to remove any unbound avidin-enzyme reagent, a substrate solution is added to the wells and color develops in proportion to the amount of MYOC bound in the initial step. The color development is stopped and the intensity of the color is measured.

Note: This product is for in vitro research use only