Pig mucin-5 subtype AC (MUC5AC) ELISA Kit

Catalog No: #EK9076

Package Size: #EK9076-1 48T #EK9076-2 96T



Orders: order@signalwayantibody.com Support: tech@signalwayantibody.com

Description

Product Name	Pig mucin-5 subtype AC (MUC5AC) ELISA Kit		
Brief Description	ELISA Kit		
Applications	ELISA		
Species Reactivity	Pig (Sus scrofa; Porcine)		
Other Names	MUC5; gastric mucin mucin 5; subtypes A and C; tracheobronchial/gastric mucin 5AC tracheobronchial mucin		
Storage	The stability of ELISA kit is determined by the loss rate of activity. The loss rate of this kit is less than 5%		
	within the expiration date under appropriate storage condition.		
	The loss rate was determined by accelerated thermal degradation test. Keep the kit at 37C for 4 and 7 days,		
	and compare O.D.values of the kit kept at 37C with that of at recommended temperature. (referring from China		
	Biological Products Standard, which was calculated by the Arrhenius equation. For ELISA kit, 4 days storage		
	at 37C can be considered as 6 months at 2 - 8C, which means 7 days at 37C equaling 12 months at 2 - 8C).		

Application Details		
Detect Range:Request Informat	on	
Sensitivity:Request Information		
Sample Type:Serum, Plasma, C	her biological fluids	
Sample Volume: 1-200 µL		
Assay Time:1-4.5h		
Detection wavelength:450 nm		

Product Description

Detection Method:SandwichTest principle:This assay employs a two-site sandwich ELISA to quantitate MUC5AC in samples. An antibody specific for MUC5AC has been pre-coated onto a microplate. Standards and samples are pipetted into the wells and anyMUC5AC present is bound by the immobilized antibody. After removing any unbound substances, a biotin-conjugated antibody specific for MUC5AC is added to the wells. After washing, Streptavidin conjugated Horseradish Peroxidase (HRP) is added to the wells. Following a wash to remove any unbound avidin-enzyme reagent, a substrate solution is added to the wells and color develops in proportion to the amount of MUC5AC bound in the initial step. The color development is stopped and the intensity of the color is measured.

Note: This product is for in vitro research use only