## Rat Mucin-13 (MUC13) ELISA Kit

Catalog No: #EK9093

Package Size: #EK9093-1 48T #EK9093-2 96T



Orders: order@signalwayantibody.com Support: tech@signalwayantibody.com

Description	
Product Name	Rat Mucin-13 (MUC13) ELISA Kit
Brief Description	ELISA Kit
Applications	ELISA
Species Reactivity	Rat (Rattus norvegicus)
Other Names	DRCC1; FLJ20063; down-regulated in colon cancer 1 mucin 13; epithelial transmembrane
Accession No.	P97881
Uniprot	P97881
Storage	The stability of ELISA kit is determined by the loss rate of activity. The loss rate of this kit is less than 5%
	within the expiration date under appropriate storage condition.
	The loss rate was determined by accelerated thermal degradation test. Keep the kit at 37C for 4 and 7 days,
	and compare O.D.values of the kit kept at 37C with that of at recommended temperature. (referring from China
	Biological Products Standard, which was calculated by the Arrhenius equation. For ELISA kit, 4 days storage
	at 37C can be considered as 6 months at 2 - 8C, which means 7 days at 37C equaling 12 months at 2 - 8C).

Detect Range:Request Information Sensitivity:Request Information Sample Type:Serum, Plasma, Other biologic Sample Volume: 1-200 µL			
Sample Type:Serum, Plasma, Other biologi			
Sample Volume: 1-200 µL	al fluids		
Assay Time:1-4.5h			
Detection wavelength:450 nm			

## **Product Description**

Detection Method:SandwichTest principle:This assay employs a two-site sandwich ELISA to quantitate MUC13 in samples. An antibody specific for MUC13 has been pre-coated onto a microplate. Standards and samples are pipetted into the wells and anyMUC13 present is bound by the immobilized antibody. After removing any unbound substances, a biotin-conjugated antibody specific for MUC13 is added to the wells. After washing, Streptavidin conjugated Horseradish Peroxidase (HRP) is added to the wells. Following a wash to remove any unbound avidin-enzyme reagent, a substrate solution is added to the wells and color develops in proportion to the amount of MUC13 bound in the initial step. The color development is stopped and the intensity of the color is measured.

Note: This product is for in vitro research use only