Mouse Tesmin (MTL5) ELISA Kit

Catalog No: #EK9246

Package Size: #EK9246-1 48T #EK9246-2 96T



Orders: order@signalwayantibody.com Support: tech@signalwayantibody.com

$\overline{}$					
	000	۱ri	nt	\sim	n
u	esc	лΠ	υı	U	ш

Product Name	Mouse Tesmin (MTL5) ELISA Kit	
Brief Description	ELISA Kit	
Applications	ELISA	
Species Reactivity	Mouse (Mus musculus)	
Other Names	CXCDC2; MTLT; TESMIN; CXC domain containing 2 metallothionein-like 5; testis-specific	
Accession No.	Q9WTJ6	
Uniprot	Q9WTJ6	
GeneID	17771;	
Storage	The stability of ELISA kit is determined by the loss rate of activity. The loss rate of this kit is less than 5%	
	within the expiration date under appropriate storage condition.	
	The loss rate was determined by accelerated thermal degradation test. Keep the kit at 37C for 4 and 7 days,	
	and compare O.D.values of the kit kept at 37C with that of at recommended temperature. (referring from China	
	Biological Draducts Standard, which was calculated by the Arrhanius equation. For ELISA kit, 4 days starage	
	Biological Products Standard, which was calculated by the Arrhenius equation. For ELISA kit, 4 days storage	

Application Details

Detect Range:Request Information		
Sensitivity:Request Information		
Sample Type:Serum, Plasma, Other biological fluids		
Sample Volume: 1-200 μL		
Assay Time:1-4.5h		
Detection wavelength:450 nm		

Product Description

Detection Method:SandwichTest principle:This assay employs a two-site sandwich ELISA to quantitate MTL5 in samples. An antibody specific for MTL5 has been pre-coated onto a microplate. Standards and samples are pipetted into the wells and anyMTL5 present is bound by the immobilized antibody. After removing any unbound substances, a biotin-conjugated antibody specific for MTL5 is added to the wells. After washing, Streptavidin conjugated Horseradish Peroxidase (HRP) is added to the wells. Following a wash to remove any unbound avidin-enzyme reagent, a substrate solution is added to the wells and color develops in proportion to the amount of MTL5 bound in the initial step. The color development is stopped and the intensity of the color is measured.

Note: This product is for in vitro research use only