Goat Cytochrome b (MT-CYB) ELISA Kit

Catalog No: #EK9260

Package Size: #EK9260-1 48T #EK9260-2 96T



Orders: order@signalwayantibody.com Support: tech@signalwayantibody.com

Description

| Product Name | Goat Cytochrome b (MT-CYB) ELISA Kit |
|--------------------|--|
| Brief Description | ELISA Kit |
| Applications | ELISA |
| Species Reactivity | Goat (Capra hircus; Caprine) |
| Other Names | MTCYB; |
| Accession No. | P24953 |
| Uniprot | P24953 |
| Storage | The stability of ELISA kit is determined by the loss rate of activity. The loss rate of this kit is less than 5% |
| | within the expiration date under appropriate storage condition. |
| | The loss rate was determined by accelerated thermal degradation test. Keep the kit at 37C for 4 and 7 days, |
| | and compare O.D.values of the kit kept at 37C with that of at recommended temperature. (referring from China |
| | Biological Products Standard, which was calculated by the Arrhenius equation. For ELISA kit, 4 days storage |
| | at 37C can be considered as 6 months at 2 - 8C, which means 7 days at 37C equaling 12 months at 2 - 8C). |
| | |

| Detect Range:Request Information Sensitivity:Request Information Sample Type:Serum, Plasma, Oth | | | | |
|---|-----------------------|--|--|--|
| | por biological fluids | | | |
| Sample Type:Serum, Plasma, Oth | or biological fluids | | | |
| | lei biological liulus | | | |
| Sample Volume: 1-200 µL | | | | |
| Assay Time:1-4.5h | | | | |
| Detection wavelength:450 nm | | | | |

Product Description

Detection Method:SandwichTest principle:This assay employs a two-site sandwich ELISA to quantitate MT-CYB in samples. An antibody specific for MT-CYB has been pre-coated onto a microplate. Standards and samples are pipetted into the wells and anyMT-CYB present is bound by the immobilized antibody. After removing any unbound substances, a biotin-conjugated antibody specific for MT-CYB is added to the wells. After washing, Streptavidin conjugated Horseradish Peroxidase (HRP) is added to the wells. Following a wash to remove any unbound avidin-enzyme reagent, a substrate solution is added to the wells and color develops in proportion to the amount of MT-CYB bound in the initial step. The color development is stopped and the intensity of the color is measured.

Note: This product is for in vitro research use only