Sheep Metallothionein-1A (MT1A) ELISA Kit

Catalog No: #EK9289



Package Size: #EK9289-1 48T #EK9289-2 96T

Orders: order@signalwayantibody.com Support: tech@signalwayantibody.com

Descri	nt	OI	n
D 00011	٣٠	٠.	

Product Name	Sheep Metallothionein-1A (MT1A) ELISA Kit	
Brief Description	ELISA Kit	
Applications	ELISA	
Species Reactivity	Sheep (Ovis aries)	
Other Names	MGC32848; MT1; MT1S; MTC; metallothionein 1A (functional) metallothionein 1S	
Accession No.	P67982	
Uniprot	P67982	
GeneID	101117184;	
Storage	The stability of ELISA kit is determined by the loss rate of activity. The loss rate of this kit is less than 5%	
	within the expiration date under appropriate storage condition.	
	The loss rate was determined by accelerated thermal degradation test. Keep the kit at 37C for 4 and 7 days,	
	and compare O.D.values of the kit kept at 37C with that of at recommended temperature. (referring from China	
	Biological Products Standard, which was calculated by the Arrhenius equation. For ELISA kit, 4 days storage	
	at 37C can be considered as 6 months at 2 - 8C, which means 7 days at 37C equaling 12 months at 2 - 8C).	

Application Details

Detect Range:Request Information
Sensitivity:Request Information
Sample Type:Serum, Plasma, Other biological fluids
Sample Volume: 1-200 μL
Assay Time:1-4.5h
Detection wavelength:450 nm

Product Description

Detection Method:SandwichTest principle:This assay employs a two-site sandwich ELISA to quantitate MT1A in samples. An antibody specific for MT1A has been pre-coated onto a microplate. Standards and samples are pipetted into the wells and anyMT1A present is bound by the immobilized antibody. After removing any unbound substances, a biotin-conjugated antibody specific for MT1A is added to the wells. After washing, Streptavidin conjugated Horseradish Peroxidase (HRP) is added to the wells. Following a wash to remove any unbound avidin-enzyme reagent, a substrate solution is added to the wells and color develops in proportion to the amount of MT1A bound in the initial step. The color development is stopped and the intensity of the color is measured.

Note: This product is for in vitro research use only