## **Product Datasheet**

## Mouse Motor neuron and pancreas homeobox protein 1 (MNX1) ELISA Kit

Catalog No: #EK9463

Package Size: #EK9463-1 48T #EK9463-2 96T



Orders: order@signalwayantibody.com Support: tech@signalwayantibody.com

Description	
Product Name	Mouse Motor neuron and pancreas homeobox protein 1 (MNX1) ELISA Kit
Brief Description	ELISA Kit
Applications	ELISA
Species Reactivity	Mouse (Mus musculus)
Other Names	HB9; HLXB9; HOXHB9; SCRA1; homeobox HB9
Accession No.	Q9QZW9
Uniprot	Q9QZW9
Storage	The stability of ELISA kit is determined by the loss rate of activity. The loss rate of this kit is less than 5%
	within the expiration date under appropriate storage condition.
	The loss rate was determined by accelerated thermal degradation test. Keep the kit at 37C for 4 and 7 days,
	and compare O.D.values of the kit kept at 37C with that of at recommended temperature. (referring from China
	Biological Products Standard, which was calculated by the Arrhenius equation. For ELISA kit, 4 days storage
	at 37C can be considered as 6 months at 2 - 8C, which means 7 days at 37C equaling 12 months at 2 - 8C).

Application Details		
Detect Range:Request Informati	n	
Sensitivity:Request Information		
Sample Type:Serum, Plasma, O	ner biological fluids	
Sample Volume: 1-200 µL		
Assay Time:1-4.5h		
Detection wavelength:450 nm		

## **Product Description**

Detection Method:SandwichTest principle:This assay employs a two-site sandwich ELISA to quantitate MNX1 in samples. An antibody specific for MNX1 has been pre-coated onto a microplate. Standards and samples are pipetted into the wells and anyMNX1 present is bound by the immobilized antibody. After removing any unbound substances, a biotin-conjugated antibody specific for MNX1 is added to the wells. After washing, Streptavidin conjugated Horseradish Peroxidase (HRP) is added to the wells. Following a wash to remove any unbound avidin-enzyme reagent, a substrate solution is added to the wells and color develops in proportion to the amount of MNX1 bound in the initial step. The color development is stopped and the intensity of the color is measured.

Note: This product is for in vitro research use only