Product Datasheet

Guinea pig Macrophage Migration Inhibitory Factor (MIF) ELISA Kit

Catalog No: #EK9586

Package Size: #EK9586-1 48T #EK9586-2 96T



Orders: order@signalwayantibody.com Support: tech@signalwayantibody.com

Description				
Product Name	Guinea pig Macrophage Migration Inhibitory Factor (MIF) ELISA Kit			
Brief Description	ELISA Kit			
Applications	ELISA			
Species Reactivity	Guinea pig (Cavia)			
Other Names	GIF; GLIF; MMIF; macrophage migration inhibitory factor phenylpyruvate tautomerase			
Storage	The stability of ELISA kit is determined by the loss rate of activity. The loss rate of this kit is less than 5%			
	within the expiration date under appropriate storage condition.			
	The loss rate was determined by accelerated thermal degradation test. Keep the kit at 37C for 4 and 7 days,			
	and compare O.D.values of the kit kept at 37C with that of at recommended temperature. (referring from China			
	Biological Products Standard, which was calculated by the Arrhenius equation. For ELISA kit, 4 days storage			
	at 37C can be considered as 6 months at 2 - 8C, which means 7 days at 37C equaling 12 months at 2 - 8C).			

Detect Range:Request Information Sensitivity:Request Information Sample Type:Serum, Plasma, Other biological fluids Sample Volume: 1-200 µL Assay Time:1-4.5h	Application Details			
Sample Type:Serum, Plasma, Other biological fluids Sample Volume: 1-200 µL	Detect Range:Request Information	tion		
Sample Volume: 1-200 µL	Sensitivity:Request Information			
	Sample Type:Serum, Plasma,	Other biological fluids		
Assay Time:1-4.5h	Sample Volume: 1-200 µL			
	Assay Time:1-4.5h			
Detection wavelength:450 nm	Detection wavelength:450 nm			

Product Description

Detection Method:SandwichTest principle:This assay employs a two-site sandwich ELISA to quantitate MIF in samples. An antibody specific for MIF has been pre-coated onto a microplate. Standards and samples are pipetted into the wells and anyMIF present is bound by the immobilized antibody. After removing any unbound substances, a biotin-conjugated antibody specific for MIF is added to the wells. After washing, Streptavidin conjugated Horseradish Peroxidase (HRP) is added to the wells. Following a wash to remove any unbound avidin-enzyme reagent, a substrate solution is added to the wells and color develops in proportion to the amount of MIF bound in the initial step. The color development is stopped and the intensity of the color is measured.

Note: This product is for in vitro research use only