

# Mouse Mitogen-activated protein kinase-binding protein 1 (MAPKBP1) ELISA Kit



Catalog No: #EK9906

Orders: order@signalwayantibody.com

Package Size: #EK9906-1 48T #EK9906-2 96T

Support: tech@signalwayantibody.com

## Description

Product Name	Mouse Mitogen-activated protein kinase-binding protein 1 (MAPKBP1) ELISA Kit
Brief Description	ELISA Kit
Applications	ELISA
Species Reactivity	Mouse (Mus musculus)
Other Names	MGC138851; MGC138852; mitogen activated protein kinase binding protein 1 mitogen-activated protein kinase binding protein 1-like
Accession No.	Q6NS57
Uniprot	Q6NS57
GeneID	26390;
Storage	The stability of ELISA kit is determined by the loss rate of activity. The loss rate of this kit is less than 5% within the expiration date under appropriate storage condition.  The loss rate was determined by accelerated thermal degradation test. Keep the kit at 37C for 4 and 7 days, and compare O.D.values of the kit kept at 37C with that of at recommended temperature. (referring from China Biological Products Standard, which was calculated by the Arrhenius equation. For ELISA kit, 4 days storage at 37C can be considered as 6 months at 2 - 8C, which means 7 days at 37C equaling 12 months at 2 - 8C).

## Application Details

Detect Range:Request Information

Sensitivity:Request Information

Sample Type:Serum, Plasma, Other biological fluids

Sample Volume: 1-200 µL

Assay Time:1-4.5h

Detection wavelength:450 nm

## Product Description

Detection Method:SandwichTest principle:This assay employs a two-site sandwich ELISA to quantitate MAPKBP1 in samples. An antibody specific for MAPKBP1 has been pre-coated onto a microplate. Standards and samples are pipetted into the wells and anyMAPKBP1 present is bound by the immobilized antibody. After removing any unbound substances, a biotin-conjugated antibody specific for MAPKBP1 is added to the wells. After washing, Streptavidin conjugated Horseradish Peroxidase (HRP) is added to the wells. Following a wash to remove any unbound avidin-enzyme reagent, a substrate solution is added to the wells and color develops in proportion to the amount of MAPKBP1 bound in the initial step. The color development is stopped and the intensity of the color is measured.Product Overview:MAPKBP1 is expressed at high level, 1.7 times the average gene in this release. MAPKBP1 contains 40 different introns (39 gt-ag, 1 gc-ag). Transcription produces 20 different mRNAs, 14 alternatively spliced variants and 6 unspliced forms. There are 6 probable alternative promoters, 6 non overlapping alternative last exons and 2 validated alternative polyadenylation sites (see the diagram). The mRNAs appear to differ by truncation of the 5' end, truncation of the 3' end, presence or absence of 26 cassette exons, overlapping exons with different boundaries, alternative splicing or retention of 10 introns. 1495 bp of this gene are antisense to spliced gene plargley, raising the possibility of regulated alternate expression. There are 2 articles specifically referring to this gene in PubMed. Proteins are expected to localize in nucleus.

---

Note: This product is for in vitro research use only