Product Datasheet

Bovine Ficolin-1 (FCN1) ELISA Kit

Catalog No: #EK9965

Package Size: #EK9965-1 48T #EK9965-2 96T



Orders: order@signalwayantibody.com Support: tech@signalwayantibody.com

_					
	20	۱'n	nt	חו	n
$\mathbf{\nu}$	esc	/I II	νι	ıv	ш

Product Name	Bovine Ficolin-1 (FCN1) ELISA Kit	
Brief Description	ELISA Kit	
Applications	ELISA	
Species Reactivity	Bovine (Bos taurus; Cattle)	
Other Names	FCNM; ficolin (collagen/fibrinogen domain-containing) 1 ficolin 1	
Accession No.	O00602	
Uniprot	O00602	
GeneID	2219;	
Storage	The stability of ELISA kit is determined by the loss rate of activity. The loss rate of this kit is less than 5%	
	within the expiration date under appropriate storage condition.	
	The loss rate was determined by accelerated thermal degradation test. Keep the kit at 37C for 4 and 7 days,	
	and compare O.D.values of the kit kept at 37C with that of at recommended temperature. (referring from China	
	Biological Products Standard, which was calculated by the Arrhenius equation. For ELISA kit, 4 days storage	
	at 37C can be considered as 6 months at 2 - 8C, which means 7 days at 37C equaling 12 months at 2 - 8C).	

Application Details

Detect Range:1.56-100 ng/mL

Sensitivity:0.78 ng/mL

Sample Type:Serum, Plasma, Other biological fluids

Sample Volume: 1-200 µL

Assay Time:1-4.5h

Detection wavelength:450 nm

Product Description

Detection Method:SandwichTest principle:This assay employs a two-site sandwich ELISA to quantitate FCN1 in samples. An antibody specific for FCN1 has been pre-coated onto a microplate. Standards and samples are pipetted into the wells and anyFCN1 present is bound by the immobilized antibody. After removing any unbound substances, a biotin-conjugated antibody specific for FCN1 is added to the wells. After washing, Streptavidin conjugated Horseradish Peroxidase (HRP) is added to the wells. Following a wash to remove any unbound avidin-enzyme reagent, a substrate solution is added to the wells and color develops in proportion to the amount of FCN1 bound in the initial step. The color development is stopped and the intensity of the color is measured. Product Overview: The ficolin family of proteins are characterized by the presence of a leader peptide, a short N-terminal segment, followed by a collagen-like region, and a C-terminal fibrinogen-like domain. The collagen-like and the fibrinogen-like domains are also found separately in other proteins such as complement protein C1q, C-type lectins known as collectins, and tenascins. However, all these proteins recognize different targets, and are functionally distinct. Ficolin 1 encoded by FCN1 is predominantly expressed in the peripheral blood leukocytes, and has been postulated to function as a plasma protein with elastin-binding activity.

Note: This product is for in vitro research use only