## Cat Interleukin 6 (IL-6) ELISA Kit

Catalog No: #EK10270

Package Size: #EK10270-1 48T #EK10270-2 96T



Orders: order@signalwayantibody.com Support: tech@signalwayantibody.com

ELISA Kit ELISA Cat (Felis catus,Feline) BSF2; HGF; HSF; IFNB2; IL-6; B cell stimulatory factor-2 B-cell differentiation factor CTL differentiation factor OTTHUMP00000158544 hybridoma growth factor interleukin 6 interleukin BSF-2 P41683 P41683
Cat (Felis catus,Feline) BSF2; HGF; HSF; IFNB2; IL-6; B cell stimulatory factor-2 B-cell differentiation factor CTL differentiation factor OTTHUMP00000158544 hybridoma growth factor interleukin 6 interleukin BSF-2 P41683 P41683
BSF2; HGF; HSF; IFNB2; IL-6; B cell stimulatory factor-2 B-cell differentiation factor CTL differentiation factor OTTHUMP00000158544 hybridoma growth factor interleukin 6 interleukin BSF-2 P41683 P41683
factor OTTHUMP00000158544 hybridoma growth factor interleukin 6 interleukin BSF-2 P41683 P41683
P41683
P41683
403687
43007,
The stability of ELISA kit is determined by the loss rate of activity. The loss rate of this kit is less than 5% within the expiration date under appropriate storage condition. The loss rate was determined by accelerated thermal degradation test. Keep the kit at 37C for 4 and 7 days, and compare O.D.values of the kit kept at 37C with that of at recommended temperature. (referring from China Biological Products Standard, which was calculated by the Arrhenius equation. For ELISA kit, 4 days storage
т а

Application Details	
Detect Range:6.25-400 pg/mL	
Sensitivity:4.5 pg/mL	
Sample Type:Serum, Plasma,	Other biological fluids
Sample Volume: 1-200 µL	
Assay Time:1-4.5h	
Detection wavelength:450 nm	

## **Product Description**

Detection Method:SandwichTest principle:This assay employs a two-site sandwich ELISA to quantitate IL6 in samples. An antibody specific for IL6 has been pre-coated onto a microplate. Standards and samples are pipetted into the wells and anyIL6 present is bound by the immobilized antibody. After removing any unbound substances, a biotin-conjugated antibody specific for IL6 is added to the wells. After washing, Streptavidin conjugated Horseradish Peroxidase (HRP) is added to the wells. Following a wash to remove any unbound avidin-enzyme reagent, a substrate solution is added to the wells and color develops in proportion to the amount of IL6 bound in the initial step. The color development is stopped and the intensity of the color is measured.

Note: This product is for in vitro research use only