Pig Interleukin-1 beta (IL1B) ELISA Kit

Catalog No: #EK10291

Package Size: #EK10291-1 48T #EK10291-2 96T



Orders: order@signalwayantibody.com Support: tech@signalwayantibody.com

Description	
Product Name	Pig Interleukin-1 beta (IL1B) ELISA Kit
Brief Description	ELISA Kit
Applications	ELISA
Species Reactivity	Pig (Sus scrofa; Porcine)
Other Names	IL-1; IL1-BETA; IL1F2; catabolin preinterleukin 1 beta pro-interleukin-1-beta
Accession No.	P26889
Uniprot	P26889
GeneID	397122;
Storage	The stability of ELISA kit is determined by the loss rate of activity. The loss rate of this kit is less than 5% within the expiration date under appropriate storage condition.
	The loss rate was determined by accelerated thermal degradation test. Keep the kit at 37C for 4 and 7 days,
	and compare O.D.values of the kit kept at 37C with that of at recommended temperature. (referring from China
	Biological Products Standard, which was calculated by the Arrhenius equation. For ELISA kit, 4 days storage
	at 37C can be considered as 6 months at 2 - 8C, which means 7 days at 37C equaling 12 months at 2 - 8C).

Detect Range:3.125-200 pg/mL Sensitivity:0.9 pg/mL Sample Type:Serum, Plasma, Other biological fluids Sample Volume: 1-200 µL Assay Time:1-4.5h Detection wavelength:450 nm	Application Details		
Sample Type:Serum, Plasma, Other biological fluids Sample Volume: 1-200 µL Assay Time:1-4.5h	Detect Range:3.125-200 pg/ml		
Sample Volume: 1-200 µL Assay Time:1-4.5h	Sensitivity:0.9 pg/mL		
Assay Time:1-4.5h	Sample Type:Serum, Plasma,	Other biological fluids	
•	Sample Volume: 1-200 µL		
Detection wavelength: 450 nm	Assay Time:1-4.5h		
	Detection wavelength:450 nm		

Product Description

Detection Method:SandwichTest principle:This assay employs a two-site sandwich ELISA to quantitate IL1B in samples. An antibody specific for IL1B has been pre-coated onto a microplate. Standards and samples are pipetted into the wells and anyIL1B present is bound by the immobilized antibody. After removing any unbound substances, a biotin-conjugated antibody specific for IL1B is added to the wells. After washing, Streptavidin conjugated Horseradish Peroxidase (HRP) is added to the wells. Following a wash to remove any unbound avidin-enzyme reagent, a substrate solution is added to the wells and color develops in proportion to the amount of IL1B bound in the initial step. The color development is stopped and the intensity of the color is measured.

Note: This product is for in vitro research use only