Human Probable G-protein coupled receptor 132 (GPR132) ELISA Kit

SAB Signalway Antibody

Catalog No: #EK10402

Package Size: #EK10402-1 48T #EK10402-2 96T

Orders: order@signalwayantibody.com Support: tech@signalwayantibody.com

Description

Product Name	Human Probable G-protein coupled receptor 132 (GPR132) ELISA Kit		
Brief Description	ELISA Kit		
Applications	ELISA		
Species Reactivity	Human (Homo sapiens)		
Other Names	G2A; MGC99642; G protein-coupled receptor G2A G2 accumulation protein OTTHUMP00000178739		
Accession No.	Q9UNW8		
Uniprot	Q9UNW8		
GeneID	29933;		
Storage	The stability of ELISA kit is determined by the loss rate of activity. The loss rate of this kit is less than 5%		
	within the expiration date under appropriate storage condition.		
	The loss rate was determined by accelerated thermal degradation test. Keep the kit at 37C for 4 and 7 days,		
	and compare O.D.values of the kit kept at 37C with that of at recommended temperature. (referring from China		
	Biological Products Standard, which was calculated by the Arrhenius equation. For ELISA kit, 4 days storage		
	at 37C can be considered as 6 months at 2 - 8C, which means 7 days at 37C equaling 12 months at 2 - 8C).		

Application Details

Detect Range:15.6-1000 pg/mL			
Sensitivity:7.5 pg/mL			
Sample Type:Serum, Plasma, Other biological fluids			
Sample Volume: 1-200 µL			
Assay Time:1-4.5h			
Detection wavelength:450 nm			

Product Description

Detection Method:SandwichTest principle:This assay employs a two-site sandwich ELISA to quantitate GPR132 in samples. An antibody specific for GPR132 has been pre-coated onto a microplate. Standards and samples are pipetted into the wells and anyGPR132 present is bound by the immobilized antibody. After removing any unbound substances, a biotin-conjugated antibody specific for GPR132 is added to the wells. After washing, Streptavidin conjugated Horseradish Peroxidase (HRP) is added to the wells. Following a wash to remove any unbound avidin-enzyme reagent, a substrate solution is added to the wells and color develops in proportion to the amount of GPR132 bound in the initial step. The color development is stopped and the intensity of the color is measured.

Note: This product is for in vitro research use only