Mouse Glutamate decarboxylase 1 (GAD1) ELISA Kit

Catalog No: #EK10429



Package Size: #EK10429-1 48T #EK10429-2 96T Orders: order@signalwayantil

Orders: order@signalwayantibody.com	
Support: tech@signalwayantibody.com	

Description			
Product Name	Mouse Glutamate decarboxylase 1 (GAD1) ELISA Kit		
Brief Description	ELISA Kit		
Applications	ELISA		
Species Reactivity	Mouse (Mus musculus)		
Other Names	FLJ45882; GAD; SCP; glutamate decarboxylase 1		
Accession No.	P48318		
Uniprot	P48318		
GeneID	14415;		
Storage	The stability of ELISA kit is determined by the loss rate of activity. The loss rate of this kit is less than 5%		
	within the expiration date under appropriate storage condition.		
	The loss rate was determined by accelerated thermal degradation test. Keep the kit at 37C for 4 and 7 days,		
	and compare O.D.values of the kit kept at 37C with that of at recommended temperature. (referring from China		
	Biological Products Standard, which was calculated by the Arrhenius equation. For ELISA kit, 4 days storage		
	at 37C can be considered as 6 months at 2 - 8C, which means 7 days at 37C equaling 12 months at 2 - 8C).		

Α .		— ''	
Λ \sim \sim	lication	しいつせつい	

Detect Range:0.78-50 ng/mL	
Sensitivity:0.4 ng/mL	
Sample Type:Serum, Plasma, Other biological fluids	
Sample Volume: 1-200 μL	
Assay Time:1-4.5h	
Detection wavelength:450 nm	

Product Description

Detection Method:SandwichTest principle:This assay employs a two-site sandwich ELISA to quantitate GAD1 in samples. An antibody specific for GAD1 has been pre-coated onto a microplate. Standards and samples are pipetted into the wells and anyGAD1 present is bound by the immobilized antibody. After removing any unbound substances, a biotin-conjugated antibody specific for GAD1 is added to the wells. After washing, Streptavidin conjugated Horseradish Peroxidase (HRP) is added to the wells. Following a wash to remove any unbound avidin-enzyme reagent, a substrate solution is added to the wells and color develops in proportion to the amount of GAD1 bound in the initial step. The color development is stopped and the intensity of the color is measured.

Note: This product is for in vitro research use only