

# Mouse Fragile X mental retardation 1 neighbor protein (FMR1NB) ELISA Kit



Catalog No: #EK10447

Orders: [order@signalwayantibody.com](mailto:order@signalwayantibody.com)

Package Size: #EK10447-1 48T #EK10447-2 96T

Support: [tech@signalwayantibody.com](mailto:tech@signalwayantibody.com)

## Description

Product Name	Mouse Fragile X mental retardation 1 neighbor protein (FMR1NB) ELISA Kit
Brief Description	ELISA Kit
Applications	ELISA
Species Reactivity	Mouse (Mus musculus)
Other Names	CT37; FLJ25736; NY-SAR-35; NYSAR35; cancer/testis antigen 37
Accession No.	Q80ZA7
Uniprot	Q80ZA7
GeneID	207854;
Storage	<p>The stability of ELISA kit is determined by the loss rate of activity. The loss rate of this kit is less than 5% within the expiration date under appropriate storage condition.</p> <p>The loss rate was determined by accelerated thermal degradation test. Keep the kit at 37C for 4 and 7 days, and compare O.D.values of the kit kept at 37C with that of at recommended temperature. (referring from China Biological Products Standard, which was calculated by the Arrhenius equation. For ELISA kit, 4 days storage at 37C can be considered as 6 months at 2 - 8C, which means 7 days at 37C equaling 12 months at 2 - 8C).</p>

## Application Details

Detect Range:15.6-1000 pg/mL

Sensitivity:7.5 pg/mL

Sample Type:Serum, Plasma, Other biological fluids

Sample Volume: 1-200 µL

Assay Time:1-4.5h

Detection wavelength:450 nm

## Product Description

Detection Method:SandwichTest principle:This assay employs a two-site sandwich ELISA to quantitate FMR1NB in samples. An antibody specific for FMR1NB has been pre-coated onto a microplate. Standards and samples are pipetted into the wells and anyFMR1NB present is bound by the immobilized antibody. After removing any unbound substances, a biotin-conjugated antibody specific for FMR1NB is added to the wells. After washing, Streptavidin conjugated Horseradish Peroxidase (HRP) is added to the wells. Following a wash to remove any unbound avidin-enzyme reagent, a substrate solution is added to the wells and color develops in proportion to the amount of FMR1NB bound in the initial step. The color development is stopped and the intensity of the color is measured.

Note: This product is for in vitro research use only