Human Ephrin-A2 (EFNA2) ELISA Kit

Catalog No: #EK10496

Package Size: #EK10496-1 48T #EK10496-2 96T



Orders: order@signalwayantibody.com Support: tech@signalwayantibody.com

| Product Name | Human Ephrin-A2 (EFNA2) ELISA Kit |
|--------------------|--|
| Brief Description | ELISA Kit |
| Applications | ELISA |
| Species Reactivity | Human (Homo sapiens) |
| Other Names | ELF-1; EPLG6; HEK7-L; LERK6; HEK7-ligand eph-related receptor tyrosine kinase ligand 6 ligand of |
| | eph-related kinase 6 |
| Accession No. | O43921 |
| Uniprot | O43921 |
| GeneID | 1943; |
| Storage | The stability of ELISA kit is determined by the loss rate of activity. The loss rate of this kit is less than 5% |
| | within the expiration date under appropriate storage condition. |
| | The loss rate was determined by accelerated thermal degradation test. Keep the kit at 37C for 4 and 7 days, |
| | and compare O.D.values of the kit kept at 37C with that of at recommended temperature. (referring from China |
| | Biological Products Standard, which was calculated by the Arrhenius equation. For ELISA kit, 4 days storage |
| | at 37C can be considered as 6 months at 2 - 8C, which means 7 days at 37C equaling 12 months at 2 - 8C). |

| Application Details | |
|-----------------------------|-------------------------|
| Detect Range:0.156-10 ng/mL | |
| Sensitivity:0.055 ng/mL | |
| Sample Type:Serum, Plasma, | Other biological fluids |
| Sample Volume: 1-200 µL | |
| Assay Time:1-4.5h | |
| Detection wavelength:450 nm | |
| | |

Product Description

Detection Method:SandwichTest principle:This assay employs a two-site sandwich ELISA to quantitate EFNA2 in samples. An antibody specific for EFNA2 has been pre-coated onto a microplate. Standards and samples are pipetted into the wells and anyEFNA2 present is bound by the immobilized antibody. After removing any unbound substances, a biotin-conjugated antibody specific for EFNA2 is added to the wells. After washing, Streptavidin conjugated Horseradish Peroxidase (HRP) is added to the wells. Following a wash to remove any unbound avidin-enzyme reagent, a substrate solution is added to the wells and color develops in proportion to the amount of EFNA2 bound in the initial step. The color development is stopped and the intensity of the color is measured.Product Overview:Ephrin-A2 is a member of the ephrin family. The protein is composed of a signal sequence, a receptor-binding region, a spacer region, and a hydrophobic region. The EPH and EPH-related receptors comprise the largest subfamily of receptor protein-tyrosine kinases and have been implicated in mediating developmental events, particularly in the nervous system. Based on their structures and sequence relationships, ephrins are divided into the ephrin-A (EFNA) class, which are anchored to the membrane by a glycosylphosphatidylinositol linkage, and the ephrin-B (EFNB) class, which are transmembrane proteins. Posttranslational modifications determine

whether this protein localizes to the nucleus or the cytoplasm.

Note: This product is for in vitro research use only