

Human Dopamine D2 receptor (D2R) ELISA Kit

Catalog No: #EK10532



Package Size: #EK10532-1 48T #EK10532-2 96T

Orders: order@signalwayantibody.com

Support: tech@signalwayantibody.com

Description

Product Name	Human Dopamine D2 receptor (D2R) ELISA Kit
Brief Description	ELISA Kit
Applications	ELISA
Species Reactivity	Human (Homo sapiens)
Other Names	D2DR; D2R; dopamine receptor D2 isoform seven transmembrane helix receptor
Accession No.	P14416
Uniprot	P14416
GeneID	1813;
Storage	<p>The stability of ELISA kit is determined by the loss rate of activity. The loss rate of this kit is less than 5% within the expiration date under appropriate storage condition.</p> <p>The loss rate was determined by accelerated thermal degradation test. Keep the kit at 37C for 4 and 7 days, and compare O.D.values of the kit kept at 37C with that of at recommended temperature. (referring from China Biological Products Standard, which was calculated by the Arrhenius equation. For ELISA kit, 4 days storage at 37C can be considered as 6 months at 2 - 8C, which means 7 days at 37C equaling 12 months at 2 - 8C).</p>

Application Details

Detect Range:0.313-20 ng/mL

Sensitivity:0.123 ng/mL

Sample Type:Serum, Plasma, Other biological fluids

Sample Volume: 1-200 µL

Assay Time:1-4.5h

Detection wavelength:450 nm

Product Description

Detection Method:SandwichTest principle:This assay employs a two-site sandwich ELISA to quantitate DRD2 in samples. An antibody specific for DRD2 has been pre-coated onto a microplate. Standards and samples are pipetted into the wells and anyDRD2 present is bound by the immobilized antibody. After removing any unbound substances, a biotin-conjugated antibody specific for DRD2 is added to the wells. After washing, Streptavidin conjugated Horseradish Peroxidase (HRP) is added to the wells. Following a wash to remove any unbound avidin-enzyme reagent, a substrate solution is added to the wells and color develops in proportion to the amount of DRD2 bound in the initial step. The color development is stopped and the intensity of the color is measured.Product Overview:D2 (DRD2). There is a short version of D2 (D2Sh) and a long version of D2 (D2Lh): The D2Sh are pre-synaptic situated, having modulatory functions (called autoreceptor, they regulate the neurotransmission by feed-back mechanisms, i.e., synthesis, storage and release of dopamine into the synaptic cleft).

The D2Lh may have the classic function of a post-synaptic receptor, i.e., keep going on the neurotransmission (excitatory or inhibitory) once blocked by a receptor antagonist or stimulated by the endogenous neurotransmitter itself or a synthetic full or partial agonist.D2-like activation is coupled to the G protein Gai, which subsequently increases phosphodiesterase activity. Phosphodiesterases break down cAMP, producing an inhibitory effect in neurons.

Note: This product is for in vitro research use only