

Rat Disks large-associated protein 3 (DLGAP3) ELISA Kit



Catalog No: #EK10609

Orders: order@signalwayantibody.com

Package Size: #EK10609-1 48T #EK10609-2 96T

Support: tech@signalwayantibody.com

Description

Product Name	Rat Disks large-associated protein 3 (DLGAP3) ELISA Kit
Brief Description	ELISA Kit
Applications	ELISA
Species Reactivity	Rat (<i>Rattus norvegicus</i>)
Other Names	DAP3; SAPAP3; OTTHUMP0000004180 PSD-95/SAP90-binding protein 3 SAP90/PSD 95 associated protein 3 SAP90/PSD-95-associated protein 3 disks large-associated protein 3
Accession No.	P97838
Uniprot	P97838
Storage	<p>The stability of ELISA kit is determined by the loss rate of activity. The loss rate of this kit is less than 5% within the expiration date under appropriate storage condition.</p> <p>The loss rate was determined by accelerated thermal degradation test. Keep the kit at 37C for 4 and 7 days, and compare O.D.values of the kit kept at 37C with that of at recommended temperature. (referring from China Biological Products Standard, which was calculated by the Arrhenius equation. For ELISA kit, 4 days storage at 37C can be considered as 6 months at 2 - 8C, which means 7 days at 37C equaling 12 months at 2 - 8C).</p>

Application Details

Detect Range:Request Information

Sensitivity:Request Information

Sample Type:Serum, Plasma, Other biological fluids

Sample Volume: 1-200 µL

Assay Time:1-4.5h

Detection wavelength:450 nm

Product Description

Detection Method:SandwichTest principle:This assay employs a two-site sandwich ELISA to quantitate DLGAP3 in samples. An antibody specific for DLGAP3 has been pre-coated onto a microplate. Standards and samples are pipetted into the wells and anyDLGAP3 present is bound by the immobilized antibody. After removing any unbound substances, a biotin-conjugated antibody specific for DLGAP3 is added to the wells. After washing, Streptavidin conjugated Horseradish Peroxidase (HRP) is added to the wells. Following a wash to remove any unbound avidin-enzyme reagent, a substrate solution is added to the wells and color develops in proportion to the amount of DLGAP3 bound in the initial step. The color development is stopped and the intensity of the color is measured.**Product Overview:**Using a segment of rat Fak (PTK2) to screen an adult human brain cDNA library, Bongiorno-Borbone et al. (2005) cloned DLGAP3, which they called SAPAP3. The deduced protein contains 977 amino acids. By Northern blot analysis, Takeuchi et al. (1997) found brain-specific expression of rat Sapap3. The rat protein localized to dendrites and cell bodies and was enriched in synaptosomal membrane fractions.By mutation analysis, Bongiorno-Borbone et al. (2005) demonstrated that the N-terminal half of human SAPAP3 interacted with a C-terminal segment of rat Fak. They also found that SAPAP3 interacted with Pyk2 (PTK2B) in in vitro pull-down assays. In rat forebrain lysates, part of Fak and Pyk2 had a similar subcellular distribution as Sapap3 and Sap90/Psd95, which supported their functional association.

Note: This product is for in vitro research use only