Monkey Beta-defensin 118 (DEFB118) ELISA Kit

Catalog No: #EK10823



Package Size: #EK10823-1 48T #EK10823-2 96T

Orders: order@signalwayantibody.com Support: tech@signalwayantibody.com

Description	
Product Name	Monkey Beta-defensin 118 (DEFB118) ELISA Kit
Brief Description	ELISA Kit
Applications	ELISA
Species Reactivity	Monkey (Simian)
Other Names	C20orf63; DEFB-18; ESC42; dJ1018D12.3; beta-defensin 118 defensin; beta 18 epididymus specific clone 42
Accession No.	Q95L10
Uniprot	Q95L10
GeneID	641438;
Storage	The stability of ELISA kit is determined by the loss rate of activity. The loss rate of this kit is less than 5%
	within the expiration date under appropriate storage condition.
	The loss rate was determined by accelerated thermal degradation test. Keep the kit at 37C for 4 and 7 days,
	and compare O.D.values of the kit kept at 37C with that of at recommended temperature. (referring from China
	Biological Products Standard, which was calculated by the Arrhenius equation. For ELISA kit, 4 days storage
	at 37C can be considered as 6 months at 2 - 8C, which means 7 days at 37C equaling 12 months at 2 - 8C).

Application Details

Detect Range:Request Information	
Sensitivity:Request Information	
Sample Type:Serum, Plasma, Other biological fluids	
Sample Volume: 1-200 μL	
Assay Time:1-4.5h	
Detection wavelength:450 nm	

Product Description

Detection Method:SandwichTest principle:This assay employs a two-site sandwich ELISA to quantitate DEFB118 in samples. An antibody specific for DEFB118 has been pre-coated onto a microplate. Standards and samples are pipetted into the wells and anyDEFB118 present is bound by the immobilized antibody. After removing any unbound substances, a biotin-conjugated antibody specific for DEFB118 is added to the wells. After washing, Streptavidin conjugated Horseradish Peroxidase (HRP) is added to the wells. Following a wash to remove any unbound avidin-enzyme reagent, a substrate solution is added to the wells and color develops in proportion to the amount of DEFB118 bound in the initial step. The color development is stopped and the intensity of the color is measured.

Note: This product is for in vitro research use only