## Rat Parafibromin (CDC73) ELISA Kit

Catalog No: #EK11096

SAB Signalway Antibody

Package Size: #EK11096-1 48T #EK11096-2 96T

Orders: order@signalwayantibody.com Support: tech@signalwayantibody.com

	000	¬rı	n	TI.	$\smallfrown$	n
u	eso	лΠ	U	u	U	ш

Product Name	Rat Parafibromin (CDC73) ELISA Kit			
Brief Description	ELISA Kit			
Applications	ELISA			
Species Reactivity	Rat (Rattus norvegicus)			
Other Names	C1orf28; FLJ23316; HPTJT; HRPT2; hyperparathyroidism 2 protein parafibromin			
Accession No.	Q4V8C8			
Uniprot	Q4V8C8			
GeneID	304832;			
Storage	The stability of ELISA kit is determined by the loss rate of activity. The loss rate of this kit is less than 5%			
	within the expiration date under appropriate storage condition.			
	The loss rate was determined by accelerated thermal degradation test. Keep the kit at 37C for 4 and 7 days,			
	and compare O.D.values of the kit kept at 37C with that of at recommended temperature. (referring from China			
	Biological Products Standard, which was calculated by the Arrhenius equation. For ELISA kit, 4 days storage			
	at 37C can be considered as 6 months at 2 - 8C, which means 7 days at 37C equaling 12 months at 2 - 8C).			

## **Application Details**

Detect Range:Request Information	
Sensitivity:Request Information	
Sample Type:Serum, Plasma, Other biological fluids	
Sample Volume: 1-200 μL	
Assay Time:1-4.5h	
Detection wavelength:450 nm	

## **Product Description**

Detection Method:SandwichTest principle:This assay employs a two-site sandwich ELISA to quantitate CDC73 in samples. An antibody specific for CDC73 has been pre-coated onto a microplate. Standards and samples are pipetted into the wells and anyCDC73 present is bound by the immobilized antibody. After removing any unbound substances, a biotin-conjugated antibody specific for CDC73 is added to the wells. After washing, Streptavidin conjugated Horseradish Peroxidase (HRP) is added to the wells. Following a wash to remove any unbound avidin-enzyme reagent, a substrate solution is added to the wells and color develops in proportion to the amount of CDC73 bound in the initial step. The color development is stopped and the intensity of the color is measured.

Note: This product is for in vitro research use only