Mouse CD81 antigen (CD81) ELISA Kit

Catalog No: #EK11106



Orders: order@signalwayantibody.com Support: tech@signalwayantibody.com

Package Size: #EK11106-1 48T #EK11106-2 96T

Product Name	Mouse CD81 antigen (CD81) ELISA Kit
Brief Description	ELISA Kit
Applications	ELISA
Species Reactivity	Mouse (Mus musculus)
Other Names	S5.7; TAPA1; TSPAN28; 26 kDa cell surface protein TAPA-1 CD81 antigen CD81 antigen (target of
	antiproliferative antibody 1) target of antiproliferative antibody 1
Accession No.	P35762
Uniprot	P35762
GeneID	12520;
Storage	The stability of ELISA kit is determined by the loss rate of activity. The loss rate of this kit is less than 5%
	within the expiration date under appropriate storage condition.
	The loss rate was determined by accelerated thermal degradation test. Keep the kit at 37C for 4 and 7 days,
	and compare O.D.values of the kit kept at 37C with that of at recommended temperature. (referring from China
	Biological Products Standard, which was calculated by the Arrhenius equation. For ELISA kit, 4 days storage
	at 37C can be considered as 6 months at 2 - 8C, which means 7 days at 37C equaling 12 months at 2 - 8C).

Application Details

Detect Range:0.31-20 ng/mL		
Sensitivity:0.156 ng/mL		
Sample Type:Serum, Plasma, Other biological fluids		
Sample Volume: 1-200 μL		
Assay Time:1-4.5h		
Detection wavelength:450 nm		

Product Description

Detection Method:SandwichTest principle:This assay employs a two-site sandwich ELISA to quantitate CD81 in samples. An antibody specific for CD81 has been pre-coated onto a microplate. Standards and samples are pipetted into the wells and anyCD81 present is bound by the immobilized antibody. After removing any unbound substances, a biotin-conjugated antibody specific for CD81 is added to the wells. After washing, Streptavidin conjugated Horseradish Peroxidase (HRP) is added to the wells. Following a wash to remove any unbound avidin-enzyme reagent, a substrate solution is added to the wells and color develops in proportion to the amount of CD81 bound in the initial step. The color development is stopped and the intensity of the color is measured.

Note: This product is for in vitro research use only