Horse Cholinesterase (BCHE) ELISA Kit

Catalog No: #EK11136

Package Size: #EK11136-1 48T #EK11136-2 96T



Orders: order@signalwayantibody.com Support: tech@signalwayantibody.com

Description

Product Name	Horse Cholinesterase (BCHE) ELISA Kit
Brief Description	ELISA Kit
Applications	ELISA
Species Reactivity	Horse (Equus caballus; Equine)
Other Names	CHE1; E1;
Accession No.	P81908
Uniprot	P81908
Storage	The stability of ELISA kit is determined by the loss rate of activity. The loss rate of this kit is less than 5%
	within the expiration date under appropriate storage condition.
	The loss rate was determined by accelerated thermal degradation test. Keep the kit at 37C for 4 and 7 days,
	and compare O.D.values of the kit kept at 37C with that of at recommended temperature. (referring from China
	Biological Products Standard, which was calculated by the Arrhenius equation. For ELISA kit, 4 days storage
	at 37C can be considered as 6 months at 2 - 8C, which means 7 days at 37C equaling 12 months at 2 - 8C).

Detect Range:3.125-200 ng/mL Sensitivity:1.25 ng/mL Sample Type:Serum, Plasma, Other biological fluids Sample Volume: 1-200 µL Assay Time:1-4.5h Detection wavelength:450 nm	pplication Details
Sample Type:Serum, Plasma, Other biological fluids Sample Volume: 1-200 µL Assay Time:1-4.5h	etect Range:3.125-200 ng/ml
Sample Volume: 1-200 µL Assay Time:1-4.5h	ensitivity:1.25 ng/mL
Assay Time:1-4.5h	ample Type:Serum, Plasma,
	ample Volume: 1-200 µL
Detection wavelength:450 nm	ssay Time:1-4.5h
	etection wavelength:450 nm

Product Description

Detection Method:SandwichTest principle:This assay employs a two-site sandwich ELISA to quantitate BCHE in samples. An antibody specific for BCHE has been pre-coated onto a microplate. Standards and samples are pipetted into the wells and anyBCHE present is bound by the immobilized antibody. After removing any unbound substances, a biotin-conjugated antibody specific for BCHE is added to the wells. After washing, Streptavidin conjugated Horseradish Peroxidase (HRP) is added to the wells. Following a wash to remove any unbound avidin-enzyme reagent, a substrate solution is added to the wells and color develops in proportion to the amount of BCHE bound in the initial step. The color development is stopped and the intensity of the color is measured.

Note: This product is for in vitro research use only