

Human Fibronectin type-III domain-containing protein C4orf31 (C4orf31) ELISA Kit



Catalog No: #EK11323

Orders: order@signalwayantibody.com

Package Size: #EK11323-1 48T #EK11323-2 96T

Support: tech@signalwayantibody.com

Description

Product Name	Human Fibronectin type-III domain-containing protein C4orf31 (C4orf31) ELISA Kit
Brief Description	ELISA Kit
Applications	ELISA
Species Reactivity	Human (Homo sapiens)
Other Names	FLJ23191; hypothetical protein LOC79625
Accession No.	Q8TB73
Uniprot	Q8TB73
GeneID	79625;
Storage	<p>The stability of ELISA kit is determined by the loss rate of activity. The loss rate of this kit is less than 5% within the expiration date under appropriate storage condition.</p> <p>The loss rate was determined by accelerated thermal degradation test. Keep the kit at 37C for 4 and 7 days, and compare O.D.values of the kit kept at 37C with that of at recommended temperature. (referring from China Biological Products Standard, which was calculated by the Arrhenius equation. For ELISA kit, 4 days storage at 37C can be considered as 6 months at 2 - 8C, which means 7 days at 37C equaling 12 months at 2 - 8C).</p>

Application Details

Detect Range:15.6-1000 pg/mL

Sensitivity:5.9 pg/mL

Sample Type:Serum, Plasma, Other biological fluids

Sample Volume: 1-200 µL

Assay Time:1-4.5h

Detection wavelength:450 nm

Product Description

Detection Method:SandwichTest principle:This assay employs a two-site sandwich ELISA to quantitate C4orf31 in samples. An antibody specific for C4orf31 has been pre-coated onto a microplate. Standards and samples are pipetted into the wells and anyC4orf31 present is bound by the immobilized antibody. After removing any unbound substances, a biotin-conjugated antibody specific for C4orf31 is added to the wells. After washing, Streptavidin conjugated Horseradish Peroxidase (HRP) is added to the wells. Following a wash to remove any unbound avidin-enzyme reagent, a substrate solution is added to the wells and color develops in proportion to the amount of C4orf31 bound in the initial step. The color development is stopped and the intensity of the color is measured.**Product Overview:**NDNF is evolutionarily well conserved, being present in invertebrate animals. Multiple cysteines are conserved between species and secondary structure prediction shows that NDNF is mainly composed of beta-strands.NDNF is a glycosylated, disulfide-bonded secretory protein that contains a fibronectin type III domain. NDNF promoted migration and growth and elicited neurite outgrowth of mouse hippocampal neurons in culture. NDNF also protected cultured hippocampal neurons against excitotoxicity and amyloid beta-peptide toxicity. NDNF was exclusively expressed in the brain and spinal cord. NDNF was expressed by neurons and not by astrocytes. NDNF expression was observed in the neurons during development.NDNF is a novel neurotrophic factor derived from neurons that may be useful in the treatment of neuronal degeneration diseases and nerve injuries.

Note: This product is for in vitro research use only