

Mouse B cell activation factor (BAFF) ELISA Kit

Catalog No: #EK11357



Package Size: #EK11357-1 48T #EK11357-2 96T

Orders: order@signalwayantibody.com

Support: tech@signalwayantibody.com

Description

Product Name	Mouse B cell activation factor (BAFF) ELISA Kit
Brief Description	ELISA Kit
Applications	ELISA
Species Reactivity	Mouse (Mus musculus)
Other Names	BAFF; BLYS; CD257; DTL; TALL-1; TALL1; THANK; TNFSF20; ZTNF4; ApoL related ligand TALL-1 B-cell activating factor B-lymphocyte stimulator TNF and ApoL-related leukocyte expressed ligand 1 TNF homolo
Accession No.	Q9WU72
Uniprot	Q9WU72
GenID	24099;
Storage	The stability of ELISA kit is determined by the loss rate of activity. The loss rate of this kit is less than 5% within the expiration date under appropriate storage condition. The loss rate was determined by accelerated thermal degradation test. Keep the kit at 37C for 4 and 7 days, and compare O.D.values of the kit kept at 37C with that of at recommended temperature. (referring from China Biological Products Standard, which was calculated by the Arrhenius equation. For ELISA kit, 4 days storage at 37C can be considered as 6 months at 2 - 8C, which means 7 days at 37C equaling 12 months at 2 - 8C).

Application Details

Detect Range:12.5-800 pg/mL

Sensitivity:5.2 pg/mL

Sample Type:Serum, Plasma, Other biological fluids

Sample Volume: 1-200 µL

Assay Time:1-4.5h

Detection wavelength:450 nm

Product Description

Detection Method:SandwichTest principle:This assay employs a two-site sandwich ELISA to quantitate TNFSF13B in samples. An antibody specific for TNFSF13B has been pre-coated onto a microplate. Standards and samples are pipetted into the wells and anyTNFSF13B present is bound by the immobilized antibody. After removing any unbound substances, a biotin-conjugated antibody specific for TNFSF13B is added to the wells. After washing, Streptavidin conjugated Horseradish Peroxidase (HRP) is added to the wells. Following a wash to remove any unbound avidin-enzyme reagent, a substrate solution is added to the wells and color develops in proportion to the amount of TNFSF13B bound in the initial step. The color development is stopped and the intensity of the color is measured.**Product Overview:**B-cell activating factor (BAFF) is the tumor necrosis factor ligand superfamily member 13B (TNFLSF13B), also known as B Lymphocyte Stimulator (BLyS) and TNF- and APOL-related leukocyte expressed ligand (TALL-1) and the Dendritic cell-derived TNF-like molecule (CD257 antigen; cluster of differentiation 257). BAFF is a 285-amino acid long peptide glycoprotein which undergoes glycosylation at residue 124. It is expressed as transmembrane protein on various cell types including monocytes, dendritic cells and bone marrow stromal cells. The transmembrane form can be cleaved from the membrane, generating a soluble protein fragment. BAFF is the natural ligand of three unusual tumor necrosis factor receptors named BAFF-R, TACI, and BCMA, all of which have differing binding affinities for it.

Note: This product is for in vitro research use only