

# Mouse Hepatocyte growth factor activator (HGFC) ELISA Kit



Catalog No: #EK11573

Orders: order@signalwayantibody.com

Package Size: #EK11573-1 48T #EK11573-2 96T

Support: tech@signalwayantibody.com

## Description

Product Name	Mouse Hepatocyte growth factor activator (HGFC) ELISA Kit
Brief Description	ELISA Kit
Applications	ELISA
Species Reactivity	Mouse ( <i>Mus musculus</i> )
Other Names	RP11-529E10.2; HGFA; MGC138395; MGC138397;
Accession No.	Q9R098
Uniprot	Q9R098
GeneID	54426;
Storage	<p>The stability of ELISA kit is determined by the loss rate of activity. The loss rate of this kit is less than 5% within the expiration date under appropriate storage condition.</p> <p>The loss rate was determined by accelerated thermal degradation test. Keep the kit at 37C for 4 and 7 days, and compare O.D.values of the kit kept at 37C with that of at recommended temperature. (referring from China Biological Products Standard, which was calculated by the Arrhenius equation. For ELISA kit, 4 days storage at 37C can be considered as 6 months at 2 - 8C, which means 7 days at 37C equaling 12 months at 2 - 8C).</p>

## Application Details

Detect Range:31.25-2000 pg/mL

Sensitivity:12.3 pg/mL

Sample Type:Serum, Plasma, Other biological fluids

Sample Volume: 1-200 µL

Assay Time:1-4.5h

Detection wavelength:450 nm

## Product Description

**Detection Method:**SandwichTest principle:This assay employs a two-site sandwich ELISA to quantitate HGFC in samples. An antibody specific for HGFC has been pre-coated onto a microplate. Standards and samples are pipetted into the wells and anyHGFC present is bound by the immobilized antibody. After removing any unbound substances, a biotin-conjugated antibody specific for HGFC is added to the wells. After washing, Streptavidin conjugated Horseradish Peroxidase (HRP) is added to the wells. Following a wash to remove any unbound avidin-enzyme reagent, a substrate solution is added to the wells and color develops in proportion to the amount of HGFC bound in the initial step. The color development is stopped and the intensity of the color is measured.**Product Overview:**HGFC belongs to peptidase family S1. It is first synthesized as an inactive single-chain precursor before being activated to a heterodimeric form by endoproteolytic processing. It acts as serine protease that converts hepatocyte growth factor to the active form. Miyazawa et al. (1993) determined the partial amino acid sequence of the homologous human serine protease and used it to design degenerate primers to screen a human liver cDNA library. They identified a 655-amino acid inactive precursor protein, designated hepatocyte growth factor activator (HGFC), with a calculated molecular mass of 70,681 Da. They determined that the active protein purified from serum is derived from the C terminus of the precursor by proteolytic cleavage of the bonds between arg372-val373 and arg407-ile408.

---

Note: This product is for in vitro research use only