

## Human Protein AMAC1 (AMAC1) ELISA Kit

Catalog No: #EK11684



Package Size: #EK11684-1 48T #EK11684-2 96T

Orders: [order@signalwayantibody.com](mailto:order@signalwayantibody.com)Support: [tech@signalwayantibody.com](mailto:tech@signalwayantibody.com)

## Description

Product Name	Human Protein AMAC1 (AMAC1) ELISA Kit
Brief Description	ELISA Kit
Applications	ELISA
Species Reactivity	Human (Homo sapiens)
Other Names	FLJ40154; TMEM21A; transmembrane protein 21A
Accession No.	Q8N808
Uniprot	Q8N808
GeneID	146861;
Storage	<p>The stability of ELISA kit is determined by the loss rate of activity. The loss rate of this kit is less than 5% within the expiration date under appropriate storage condition.</p> <p>The loss rate was determined by accelerated thermal degradation test. Keep the kit at 37C for 4 and 7 days, and compare O.D.values of the kit kept at 37C with that of at recommended temperature. (referring from China Biological Products Standard, which was calculated by the Arrhenius equation. For ELISA kit, 4 days storage at 37C can be considered as 6 months at 2 - 8C, which means 7 days at 37C equaling 12 months at 2 - 8C).</p>

## Application Details

Detect Range:78.1-5000 pg/mL

Sensitivity:734 pg/mL

Sample Type:Serum, Plasma, Other biological fluids

Sample Volume: 1-200 µL

Assay Time:1-4.5h

Detection wavelength:450 nm

## Product Description

**Detection Method:**SandwichTest principle:This assay employs a two-site sandwich ELISA to quantitate AMAC1 in samples. An antibody specific for AMAC1 has been pre-coated onto a microplate. Standards and samples are pipetted into the wells and anyAMAC1 present is bound by the immobilized antibody. After removing any unbound substances, a biotin-conjugated antibody specific for AMAC1 is added to the wells. After washing, Streptavidin conjugated Horseradish Peroxidase (HRP) is added to the wells. Following a wash to remove any unbound avidin-enzyme reagent, a substrate solution is added to the wells and color develops in proportion to the amount of AMAC1 bound in the initial step. The color development is stopped and the intensity of the color is measured.**Product Overview:**This gene is one of several Cys-Cys (CC) cytokine genes clustered on the q arm of chromosome 19. Cytokines are a family of secreted proteins involved in immunoregulatory and inflammatory processes. The CC cytokines are proteins characterized by two adjace???

Note: This product is for in vitro research use only