## Human Rabies virus IgM antibody (RV-Ab-IgM) ELISA Kit

SAB Signalway Antibody

Catalog No: #EK11785

Package Size: #EK11785-1 48T #EK11785-2 96T

Orders: order@signalwayantibody.com Support: tech@signalwayantibody.com

## Description

| Product Name       | Human Rabies virus IgM antibody (RV-Ab-IgM) ELISA Kit  |
|--------------------|--|
| Brief Description  | ELISA Kit  |
| Applications       | ELISA  |
| Species Reactivity | Human (Homo sapiens)   |
| Storage            | The stability of ELISA kit is determined by the loss rate of activity. The loss rate of this kit is less than 5% |
|                    | within the expiration date under appropriate storage condition.  |
|                    | The loss rate was determined by accelerated thermal degradation test. Keep the kit at 37C for 4 and 7 days,      |
|                    | and compare O.D.values of the kit kept at 37C with that of at recommended temperature. (referring from China     |
|                    | Biological Products Standard, which was calculated by the Arrhenius equation. For ELISA kit, 4 days storage      |
|                    | at 37C can be considered as 6 months at 2 - 8C, which means 7 days at 37C equaling 12 months at 2 - 8C).         |

## **Application Details**

| Detect Range:Detection antibody                    |
|--|
| Sensitivity:Request Information                    |
| Sample Type:Serum, Plasma, Other biological fluids |
| Sample Volume: 1-200 μL                            |
| Assay Time:1-4.5h                                  |
| Detection wavelength:450 nm                        |

## **Product Description**

Detection Method:Competitive ELISATest principle:This assay employs the competitive enzyme immunoassay technique. The microtiter plate provided in this kit has been pre-coated with an antibody specific to RV-Ab-IgM. Standards or samples are then added to the appropriate microtiter plate wells with a Horseradish Peroxidase (HRP)-conjugated RV-Ab-IgM and incubated. The competitive inhibition reaction is launched between with HRP labeled RV-Ab-IgM and unlabeled RV-Ab-IgM with the antibody. A substrate solution is added to the wells and the color develops in opposite to the amount of RV-Ab-IgM in the sample. The color development is stopped and the intensity of the color is measured.

Note: This product is for in vitro research use only