

# Mouse Anti-cardiolipin antibody IgM (ACA-IgM) ELISA Kit



Catalog No: #EK11789

Orders: [order@signalwayantibody.com](mailto:order@signalwayantibody.com)

Package Size: #EK11789-1 48T #EK11789-2 96T

Support: [tech@signalwayantibody.com](mailto:tech@signalwayantibody.com)

## Description

Product Name	Mouse Anti-cardiolipin antibody IgM (ACA-IgM) ELISA Kit
Brief Description	ELISA Kit
Applications	ELISA
Species Reactivity	Mouse ( <i>Mus musculus</i> )
Storage	<p>The stability of ELISA kit is determined by the loss rate of activity. The loss rate of this kit is less than 5% within the expiration date under appropriate storage condition.</p> <p>The loss rate was determined by accelerated thermal degradation test. Keep the kit at 37C for 4 and 7 days, and compare O.D.values of the kit kept at 37C with that of at recommended temperature. (referring from China Biological Products Standard, which was calculated by the Arrhenius equation. For ELISA kit, 4 days storage at 37C can be considered as 6 months at 2 - 8C, which means 7 days at 37C equaling 12 months at 2 - 8C).</p>

## Application Details

Detect Range:12.35-1000 ng/mL

Sensitivity:1.6 ng/mL

Sample Type:Serum, Plasma, Other biological fluids

Sample Volume: 1-200 µL

Assay Time:1-4.5h

Detection wavelength:450 nm

## Product Description

Detection Method:Competitive ELISATest principle:This assay employs the competitive enzyme immunoassay technique. The microtiter plate provided in this kit has been pre-coated with an antibody specific to ACA-IgM. Standards or samples are then added to the appropriate microtiter plate wells with a Horseradish Peroxidase (HRP)-conjugated ACA-IgM and incubated. The competitive inhibition reaction is launched between with HRP labeled ACA-IgM and unlabeled ACA-IgM with the antibody. A substrate solution is added to the wells and the color develops in opposite to the amount of ACA-IgM in the sample. The color development is stopped and the intensity of the color is measured.Product Overview:Cardiolipin is an important component of the inner mitochondrial membrane, where it constitutes about 22% of the total lipid composition. The name cardiolipin is derived from the fact that it was first found in animal hearts. It was first isolated from????□

Note: This product is for in vitro research use only