

# Human Tumor necrosis factor $\alpha$ IgG antibody (TNFA-Ab-IgG) ELISA Kit



Catalog No: #EK11946

Orders: order@signalwayantibody.com

Package Size: #EK11946-1 48T #EK11946-2 96T

Support: tech@signalwayantibody.com

## Description

Product Name	Human Tumor necrosis factor $\alpha$ IgG antibody (TNFA-Ab-IgG) ELISA Kit
Brief Description	ELISA Kit
Applications	ELISA
Species Reactivity	Human (Homo sapiens)
Other Names	antibody for DADB-70P7.1; DIF; TNF-alpha; TNFA; TNFSF2; APC1 protein TNF superfamily; member 2 TNF; macrophage-derived TNF; monocyte-derived cachectin tumor necrosis factor alpha
Storage	<p>The stability of ELISA kit is determined by the loss rate of activity. The loss rate of this kit is less than 5% within the expiration date under appropriate storage condition.</p> <p>The loss rate was determined by accelerated thermal degradation test. Keep the kit at 37C for 4 and 7 days, and compare O.D.values of the kit kept at 37C with that of at recommended temperature. (referring from China Biological Products Standard, which was calculated by the Arrhenius equation. For ELISA kit, 4 days storage at 37C can be considered as 6 months at 2 - 8C, which means 7 days at 37C equaling 12 months at 2 - 8C).</p>

## Application Details

Detect Range:15.6-1000 pg/mL

Sensitivity:6.1 pg/mL

Sample Type:Serum, Plasma, Other biological fluids

Sample Volume: 1-200  $\mu$ L

Assay Time:1-4.5h

Detection wavelength:450 nm

## Product Description

**Detection Method:**Sandwich**Test principle:**This assay employs a two-site sandwich ELISA to quantitate TNFA-Ab-IgG in samples. An antibody specific for TNFA-Ab-IgG has been pre-coated onto a microplate. Standards and samples are pipetted into the wells and anyTNFA-Ab-IgG present is bound by the immobilized antibody. After removing any unbound substances, a biotin-conjugated antibody specific for TNFA-Ab-IgG is added to the wells. After washing, Streptavidin conjugated Horseradish Peroxidase (HRP) is added to the wells. Following a wash to remove any unbound avidin-enzyme reagent, a substrate solution is added to the wells and color develops in proportion to the amount of TNFA-Ab-IgG bound in the initial step. The color development is stopped and the intensity of the color is measured.**Product Overview:**TNF $\alpha$ ? is synthesized as a 26 kDa, type II transmembrane protein that is 233 amino acids in length. It contains a 30 amino acid (aa) cytoplasmic domain, a 26 aa transmembrane segment, and a 177 aa extracellular region. TNF $\alpha$  is assembled intracellularly to form a transmembrane, non-covalently-linked homotrimeric protein. The 157 aa residue soluble form of TNF $\alpha$  (sTNF- $\alpha$ is released from the C-terminus of the transmembrane protein through the activity of TNF $\alpha$ -converting enzyme (TACE), a membrane -bound disintegrin metalloproteinase. Rat cells known to express TNF- $\alpha$ include B cells, colonic columnar epithelial cells, NK and CD3 CD56 hepatic natural T cells, macrophages, monocytes and monocyte-derived dendritic cells, CD4 and CD8 T cells, mast cells, neutrophils, keratinocytes, plasma cells, and adipocytes.

---

Note: This product is for in vitro research use only