

## Human Complement fragment 4a (C4a) ELISA Kit

Catalog No: #EK12084



Package Size: #EK12084-1 48T #EK12084-2 96T

Orders: order@signalwayantibody.com

Support: tech@signalwayantibody.com

## Description

Product Name	Human Complement fragment 4a (C4a) ELISA Kit
Brief Description	ELISA Kit
Applications	ELISA
Species Reactivity	Human (Homo sapiens)
Other Names	DAMC-258G8.6; C4; C4A2; C4A3; C4A4; C4A6; C4S; CO4; CPAMD2; MGC164979; RG; C4A anaphylatoxin Rodgers form of C4 acidic C4 complement component 4A
Accession No.	P0C0L4
Uniprot	P0C0L4
GeneID	720;721;
Storage	The stability of ELISA kit is determined by the loss rate of activity. The loss rate of this kit is less than 5% within the expiration date under appropriate storage condition.  The loss rate was determined by accelerated thermal degradation test. Keep the kit at 37C for 4 and 7 days, and compare O.D.values of the kit kept at 37C with that of at recommended temperature. (referring from China Biological Products Standard, which was calculated by the Arrhenius equation. For ELISA kit, 4 days storage at 37C can be considered as 6 months at 2 - 8C, which means 7 days at 37C equaling 12 months at 2 - 8C).

## Application Details

Detect Range:0.312-20 ng/mL

Sensitivity:0.112 ng/mL

Sample Type:Serum, Plasma, Other biological fluids

Sample Volume: 1-200 µL

Assay Time:1-4.5h

Detection wavelength:450 nm

## Product Description

**Detection Method:**SandwichTest principle:This assay employs a two-site sandwich ELISA to quantitate C4A in samples. An antibody specific for C4A has been pre-coated onto a microplate. Standards and samples are pipetted into the wells and anyC4A present is bound by the immobilized antibody. After removing any unbound substances, a biotin-conjugated antibody specific for C4A is added to the wells. After washing, Streptavidin conjugated Horseradish Peroxidase (HRP) is added to the wells. Following a wash to remove any unbound avidin-enzyme reagent, a substrate solution is added to the wells and color develops in proportion to the amount of C4A bound in the initial step. The color development is stopped and the intensity of the color is measured.**Product Overview:**Complement C4-A is the acidic form of complement factor 4, part of the classical activation pathway. The protein is expressed as a single chain precursor which is proteolytically cleaved into a trimer of alpha, beta, and gamma chains prior to secretion. The trimer provides a surface for interaction between the antigen-antibody complex and other complement components. The alpha chain may be cleaved to release C4 anaphylatoxin, a mediator of local inflammation. Deficiency of this protein is associated with systemic lupus erythematosus and type I diabetes mellitus. This gene localizes to the major histocompatibility complex (MHC) class III region on chromosome 6. Varying haplotypes of this gene cluster exist, such that individuals may have 1, 2, or 3 copies of this gene.

---

Note: This product is for in vitro research use only