Goat Luteinizing Hormone (LH) ELISA Kit

Catalog No: #EK12161

Package Size: #EK12161-1 48T #EK12161-2 96T



Orders: order@signalwayantibody.com Support: tech@signalwayantibody.com

Description

Product Name	Goat Luteinizing Hormone (LH) ELISA Kit		
Brief Description	ELISA Kit		
Applications	ELISA		
Species Reactivity	Goat (Capra hircus; Caprine)		
Storage	The stability of ELISA kit is determined by the loss rate of activity. The loss rate of this kit is less than 5%		
	within the expiration date under appropriate storage condition.		
	The loss rate was determined by accelerated thermal degradation test. Keep the kit at 37C for 4 and 7 days,		
	and compare O.D.values of the kit kept at 37C with that of at recommended temperature. (referring from China		
	Biological Products Standard, which was calculated by the Arrhenius equation. For ELISA kit, 4 days storage		
	at 37C can be considered as 6 months at 2 - 8C, which means 7 days at 37C equaling 12 months at 2 - 8C).		

Application Details				
Detect Range:370.37-30000 pg/mL				
Sensitivity:152.8 pg/mL				
Sample Type:Serum, Plasma, C	her biological fluids			
Sample Volume: 1-200 µL				
Assay Time:1-4.5h				
Detection wavelength:450 nm				

Product Description

Detection Method:SandwichTest principle:This assay employs a two-site sandwich ELISA to quantitate LH in samples. An antibody specific for LH has been pre-coated onto a microplate. Standards and samples are pipetted into the wells and anyLH present is bound by the immobilized antibody. After removing any unbound substances, a biotin-conjugated antibody specific for LH is added to the wells. After washing, Streptavidin conjugated Horseradish Peroxidase (HRP) is added to the wells. Following a wash to remove any unbound avidin-enzyme reagent, a substrate solution is added to the wells and color develops in proportion to the amount of LH bound in the initial step. The color development is stopped and the intensity of the color is measured.Product Overview:Luteinizing hormone (LH) is an important hormone of the reproductive system, which has found application in diagnosis and therapeutic medicine. It plays a vital role in the development and functioning of the reproductive system. Determination of LH concentration is important for detection of dysfunction of the pituitary variance axis, diagnosis of reproductive disorders, monitoring of antifertility programmes and in therapeutic preparations.LH levels are normally low during childhood and, in women, high after menopause. As LH is secreted as pulses, it is necessary to follow its concentration over a sufficient period of time to get a proper information about its blood level.During the reproductive years typical levels are between 1-20 IU/L. Physiologic high LH levels are seen during the LH surge (v.s.); typically they last 48 hours.

Note: This product is for in vitro research use only