## **Product Datasheet**

## Pig Soluble vascular endothelial growth factor receptor-2 (VEGFR-2/sFLK-1) ELISA Kit

Catalog No: #EK12217

Description

Package Size: #EK12217-1 48T #EK12217-2 96T

1 des	$(\mathbf{R})$
<i>SA</i>	B
Signalway Antibo	ody

Orders: order@signalwayantibody.com Support: tech@signalwayantibody.com

Description	
Product Name	Pig Soluble vascular endothelial growth factor receptor-2 (VEGFR-2/sFLK-1) ELISA Kit
Brief Description	ELISA Kit
Applications	ELISA
Species Reactivity	Pig (Sus scrofa; Porcine)
Storage	The stability of ELISA kit is determined by the loss rate of activity. The loss rate of this kit is less than 5%
	within the expiration date under appropriate storage condition.
	The loss rate was determined by accelerated thermal degradation test. Keep the kit at 37C for 4 and 7 days,
	and compare O.D.values of the kit kept at 37C with that of at recommended temperature. (referring from China
	Biological Products Standard, which was calculated by the Arrhenius equation. For ELISA kit, 4 days storage
	at 37C can be considered as 6 months at 2 - 8C, which means 7 days at 37C equaling 12 months at 2 - 8C).

Application Details			
Detect Range:1.56-100 ng/mL			
Sensitivity:0.62 ng/mL			
Sample Type:Serum, Plasma,	ther biological fluids		
Sample Volume: 1-200 µL			
Assay Time:1-4.5h			
Detection wavelength:450 nm			

## **Product Description**

Detection Method:SandwichTest principle:This assay employs a two-site sandwich ELISA to quantitate VEGFR-2/sFLK-1 in samples. An antibody specific for VEGFR-2/sFLK-1 has been pre-coated onto a microplate. Standards and samples are pipetted into the wells and anyVEGFR-2/sFLK-1 present is bound by the immobilized antibody. After removing any unbound substances, a biotin-conjugated antibody specific for VEGFR-2/sFLK-1 is added to the wells. After washing, Streptavidin conjugated Horseradish Peroxidase (HRP) is added to the wells. Following a wash to remove any unbound avidin-enzyme reagent, a substrate solution is added to the wells and color develops in proportion to the amount of VEGFR-2/sFLK-1 bound in the initial step. The color development is stopped and the intensity of the color is measured.

Note: This product is for in vitro research use only