

Gamma-aminobutyric acid receptor-associated protein-like 2 Polyclonal Antibody

Catalog No: #42395

Orders: order@signalwayantibody.comSupport: tech@signalwayantibody.com


Description

| | |
|-----------------------|--|
| Product Name | Gamma-aminobutyric acid receptor-associated protein-like 2 Polyclonal Antibody |
| Host Species | Rabbit |
| Clonality | Polyclonal |
| Purification | Caprylic Acid Ammonium Sulfate Precipitation purified |
| Applications | WB |
| Species Reactivity | Hu |
| Specificity | The antibody detects endogenous level of total Gamma-aminobutyric acid receptor-associated protein-like 2 polyclonal antibody. |
| Immunogen Type | protein |
| Immunogen Description | Recombinant human Gamma-aminobutyric acid receptor-associated protein-like 2 protein |
| Target Name | Gamma-aminobutyric acid receptor-associated protein-like 2 |
| Other Names | GABARAPL2 |
| Accession No. | Swiss-Prot#: P60520 |
| Uniprot | P60520 |
| GeneID | 11345; |
| Calculated MW | 13kd |
| Formulation | Preservative: 0.03% Proclin 300 Constituents: 50% Glycerol, 0.01M PBS, PH 7.4 |
| Storage | Store at -20°C |

Application Details

Western blotting: □1:500 - 1:1000

Images



Gamma-aminobutyric acid receptor-associated protein-like 2
 Antibody at 2ug/ml + 293T whole cell lysate at 20ug
 SecondaryGoat polyclonal to Rabbit IgG at 1/15000 dilution
 Predicted band size : 13kDa
 Observed band size : 60 kDa

Background

Involved in intra-Golgi traffic. Modulates intra-Golgi transport through coupling between NSF activity and SNAREs activation. It first stimulates the

ATPase activity of NSF which in turn stimulates the association with GOSR1

References

[1] "Interaction of the Unc-51-like kinase and microtubule-associated protein light chain 3 related proteins in the brain: possible role of vesicular transport in axonal elongation." Okazaki N., Yan J., Yuasa S., Ueno T., Kominami E., Masuho Y., Koga H., M

Note: This product is for in vitro research use only