

Tumor necrosis factor receptor superfamily member 10D Polyclonal Antibody

Catalog No: #42499

Orders: order@signalwayantibody.comSupport: tech@signalwayantibody.com


Description

Product Name	Tumor necrosis factor receptor superfamily member 10D Polyclonal Antibody
Host Species	Rabbit
Clonality	Polyclonal
Purification	Caprylic Acid Ammonium Sulfate Precipitation purified
Applications	IHC
Species Reactivity	Hu
Specificity	The antibody detects endogenous level of total Tumor necrosis factor receptor superfamily member 10D polyclonal antibody.
Immunogen Type	protein
Immunogen Description	Recombinant human Tumor necrosis factor receptor superfamily member 10D protein
Target Name	Tumor necrosis factor receptor superfamily member 10D
Other Names	UNQ251/PRO288, DCR2, TRAILR4, TRUNDD, TNFRSF10D, Decoy receptor 2, DcR2, TNF-related apoptosis-inducing ligand receptor 4, TRAIL receptor 4, TRAIL-R4, TRAIL receptor with a truncated death domain, CD_antigen=CD264
Accession No.	Swiss-Prot#: Q9UBN6
Uniprot	Q9UBN6
GeneID	8793;
Formulation	Preservative: 0.03% Proclin 300 Constituents: 50% Glycerol, 0.01M PBS, PH 7.4
Storage	Store at -20°C

Application Details

Immunohistochemistry: 1:20 - 1:200

Images



Immunohistochemical analysis of paraffin-embedded human kidney using #42499 at dilution of 1:100.

Background

Receptor for the cytotoxic ligand TRAIL. Contains a truncated death domain and hence is not capable of inducing apoptosis but protects against TRAIL-mediated apoptosis. Reports are contradictory with regards to its ability to induce the NF-kappa-B pathway. According to Ref.1, it cannot but according to Ref.2, it can induce the NF-kappa-B pathway.

References

[1] "A novel receptor for Apo2L/TRAIL contains a truncated death domain."Marsters S.A., Sheridan J.P., Pitti R.M., Huang A., Skubatch M., Baldwin D., Yuan J., Gurney A., Goddard A.D., Godowski P., Ashkenazi A.Curr. Biol. 7:1003-1006(1997) [2] "The nove

Note: This product is for in vitro research use only