Product Datasheet

CD44 Antibody

Catalog No: #P1035

Package Size: #P1035-1 0.1ml #P1035-2 1ml



Orders: order@signalwayantibody.com Support: tech@signalwayantibody.com

Description


| Product Name | CD44 Antibody |
|--------------------|--|
| Host Species | Mouse |
| Clonality | Monoclonal |
| Clone No. | IHC044 |
| Applications | IHC |
| Species Reactivity | Human |
| Formulation | Tris Buffer, pH 7.3 - 7.7, with 1% BSA and |
| Storage | Store at 2-8C. Do not freeze. |

Application Details

Recommended

working dilution range:1:100 - 1:200

Images



CD44 [P1035] on Breast Cancer

Product Description

Cluster of differentiation 44 (CD44) is a glycoprotein receptor for hyaluronic acid, which plays a fundamental role in cellular adhesion, stromal binding, migration, and cell-cell interactions. Studies have suggested that the CD44-hyaluronate interaction is central to tumor invasiveness. Positive staining with Anti-CD44 is implicated in a multitude of different cancer types, including breast, prostatic, renal cell, colonic, hepatocellular, and genitourinary carcinomas, as well as Non-Hodgkins Lymphoma, metastatic melanoma, gastric cancer, and some soft tissue tumors. It has also been demonstrated that there is a positive correlation between tumor progression and increased expression of CD44v, a high molecular weight CD44 isoform that has been described in epithelial cells. Given the expression of CD44 in a wide range of cancers, the most practical application of CD44 immunostaining is its use in discriminating between urothelial transitional cell carcinoma in situ from non-neoplastic changes in the urothelium.

References

Li F, et al. Cell Res. 2007; 17:3-14.

Note: This product is for in vitro research use only