TBLR1 antibody

Catalog No: #22813



Orders: order@signalwayantibody.com Support: tech@signalwayantibody.com

$\overline{}$				
	escri	n	tic	ı'n
\boldsymbol{L}	COUL	v	uv	и т

Product Name	TBLR1 antibody
Host Species	Rabbit
Clonality	Polyclonal
Purification	Purified by antigen-affinity chromatography.
Applications	WB IHC IF
Species Reactivity	Hu
Immunogen Type	Recombinant protein
Immunogen Description	Recombinant protein fragment contain a sequence corresponding to a region within amino acids 210 and 484
	of TBLR1
Target Name	TBLR1
Accession No.	Swiss-Prot:Q9BZK7Gene ID:79718
Uniprot	Q9BZK7
GeneID	79718;
Concentration	0.5mg/ml
Formulation	Supplied in 0.1M Tris-buffered saline with 10% Glycerol (pH7.0). 0.01% Thimerosal was added as a
	preservative.
Storage	Store at -20°C for long term preservation (recommended). Store at 4°C for short term use.

Application Details


Predicted MW: 56kd

Western blotting: 1:500-1:3000

Immunohistochemistry: 1:100-1:250

Immunofluorescence: 1:100-1:200

Images




Sample (30 ug of whole cell lysate)


A: A431

7.5% SDS PAGE

Primary antibody diluted at 1: 1000



Immunohistochemical analysis of paraffin-embedded Cal27 Xenograft, using TBLR1 antibody at 1: 100 dilution.



Immunofluorescence analysis of paraformaldehyde-fixed HeLa, using TBLR1 antibody at 1: 200 dilution.

Background

The protein encoded by this gene has sequence similarity with members of the WD40 repeat-containing protein family. The WD40 group is a large family of proteins, which appear to have a regulatory function. It is believed that the WD40 repeats mediate protein-protein interactions and members of the family are involved in signal transduction, RNA processing, gene regulation, vesicular trafficking, cytoskeletal assembly and may play a role in the control of cytotypic differentiation. [provided by RefSeq]

Note: This product is for in vitro research use only